

Spectral inequalities for operators on H-type groups *

J. Inglis

Imperial College London

Abstract: *In this paper the spectra of certain Schrödinger-type operators defined in the sub-elliptic setting of H-type groups are investigated. Two approaches are taken: the first one makes use of a convenient unitary transformation on the Heisenberg group, while the second proceeds via functional inequalities. The paper concludes by highlighting a major difference between the spectrum of the considered operators depending on which natural sub-Riemannian metric is chosen.*

Keywords: *H-type groups, super-Poincaré inequality, spectral gap inequality, spectrum, logarithmic Sobolev inequality*

Contents

1	Introduction	2
2	Definitions and notation	3
3	Generalisation of a classical result to the Heisenberg group	6
4	Spectral information via functional inequalities	10
4.1	Super-Poincaré inequalities	10
4.2	Potentials defined by the Carnot-Carathéodory distance	12
4.3	Potentials defined by the Kaplan distance	16
	References	26

*Supported by EPSRC EP/D05379X/1

1 Introduction

In the classical setting of \mathbb{R}^n an extensive study has been made of the spectral properties of so-called Schrödinger operators, that is operators of the form

$$\mathcal{L} = -\Delta + V \quad (1.1)$$

where Δ is the standard Laplacian on \mathbb{R}^n and V is some potential. In particular the location of the continuous and discrete parts of the spectra of such operators (if indeed they exist) is of great interest. A classical reference detailing this study is the book of M. Reed and B. Simon [21].

In this paper we pursue the same questions, but instead work in the sub-Riemannian setting of H-type groups. We consider a direct analogue of the classical operator (1.1) in such a setting, where we replace the full Laplacian with the more natural sub-Laplacian. Given an H-type group \mathbb{G} , we will thus be interested in the sub-elliptic operators

$$\mathcal{L} = -\Delta_{\mathbb{G}} + \nabla_{\mathbb{G}} U \cdot \nabla_{\mathbb{G}} \quad (1.2)$$

where $\Delta_{\mathbb{G}}$ and $\nabla_{\mathbb{G}}$ are the sub-Laplacian and sub-gradient respectively. When considered as an operator acting on $L^2(\mu_U)$ with $\mu_U \equiv Z^{-1}e^{-U}dx$, such operators are positive and self-adjoint. Our aim will be principally to find conditions on the potential U that ensure the operator (1.2) has a purely discrete spectrum, focusing on the natural case when U is given by a power of one of the intrinsic distance functions defined on the group.

Our first attempt to answer some of these questions involves the use of an appealing unitary transformation on the Heisenberg group, but it quickly becomes apparent that such an approach is limited when it comes to general H-type groups and our preferred choices for the potential U . The majority of this paper is therefore devoted to taking a functional inequality approach, similar to that taken in [7], [25] and [28], in order to overcome these limitations.

Functional inequalities on H-type groups have attracted considerable attention recently (see for example [1, 8, 9, 10, 17, 18]), since despite being relatively simple cases of sub-Riemannian settings, the structure that they possess renders traditional methods inapplicable. Indeed, the main difficulty (as described in [1]) stems from the fact that the Ricci tensor of the sub-Laplacian on an H-type group is $-\infty$, and hence the methods of Bakry and Emery (see [2]) to prove many useful inequalities can not be used.

Another major motivation for this paper is the work of Hebisch and Zegarliński ([14]), in which an effective technology was introduced to prove inequalities such as the logarithmic Sobolev inequality for measures on H-type groups of the form

$$\mu(dx) = \frac{e^{-\alpha d^p(x)}}{Z},$$

where $\alpha > 0, p > 1$ and d is Carnot-Carathéodory distance on the group. The work we present here can thus be thought of as an extension of their study to alternative inequalities that carry spectral information.

Similar questions to those addressed here have recently been investigated in [27], where general conditions for empty essential spectrum for hypoelliptic generators are put forward. However, these conditions are not easily applicable to the cases of special interest to us, i.e. when the potential U in (1.2) is defined in terms of a natural distance function on the group. This is because of the singularities possessed by such distance functions, and hence our approach is necessarily different.

The layout of this paper is as follows. After presenting the necessary definitions and notation, we start by proving a generalisation of a classical result in the setting of the Heisenberg group, which gives some conditions on the growth and smoothness of the potential U that ensure that the operator has empty essential spectrum. Since these conditions do not cover cases of particular interest to us and the methods are not easily generalised to all H-type groups, we then take a different approach, and investigate functional inequalities for the associated measures that contain spectral information. This section is split into two subsections: the first concentrates on the situation when the potential U is defined in terms of the Carnot-Carathéodory distance, while the second tries to overcome some of the extra difficulties encountered when the Carnot-Carathéodory distance is replaced by the Kaplan distance. These difficulties are perhaps rather surprising, and it is interesting that they result in operators with differently structured spectra.

2 Definitions and notation

In this section we briefly give the basic definitions and results that will be used throughout the rest of the paper. We refer the reader to [6] for a comprehensive review of this material.

Definition 2.1 (H-type group). *Let \mathfrak{g} be a finite-dimensional real Lie algebra equipped with Lie bracket $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$. Let \mathfrak{z} denote the centre of \mathfrak{g} , that is $\mathfrak{z} = \{X \in \mathfrak{g} : [X, Y] = 0 \ \forall Y \in \mathfrak{g}\}$.*

The Lie algebra \mathfrak{g} is said to be of H-type if it can be endowed with an inner product $\langle \cdot, \cdot \rangle$ such that $[\mathfrak{z}^\perp, \mathfrak{z}^\perp] = \mathfrak{z}$, and moreover, for every fixed $Z \in \mathfrak{z}$, the map $J_Z : \mathfrak{z}^\perp \rightarrow \mathfrak{z}$ defined by

$$\langle J_Z(X), Y \rangle = \langle Z, [X, Y] \rangle \quad \forall Y \in \mathfrak{z}^\perp$$

is an orthogonal map.

An H-type group is a connected and simply connected Lie group \mathbb{G} whose Lie algebra is of H-type.

In addition to this abstract definition, we have the following characterisation result (Theorem 18.2.1 of [6]):

Theorem 2.2. *\mathbb{G} is an H-type group if and only if \mathbb{G} is (isomorphic to) \mathbb{R}^{n+m} with the group law*

$$(w, z) \circ (\omega, \zeta) = \left(\begin{array}{cc} w_i + \omega_i, & i = 1, \dots, n \\ z_j + \zeta_j + \frac{1}{2} \langle U^{(j)} w, \omega \rangle, & j = 1, \dots, m \end{array} \right), \quad (2.1)$$

for $w, \omega \in \mathbb{R}^n, z, \zeta \in \mathbb{R}^m$ and where the matrices $U^{(1)}, \dots, U^{(m)}$ have the following properties:

- (1) $U^{(j)}$ is an $n \times n$ skew-symmetric and orthogonal matrix for every $j \in \{1, \dots, m\}$;
- (2) $U^{(k)} U^{(j)} + U^{(j)} U^{(k)} = 0$ for every $k, j \in \{1, \dots, m\}$ with $k \neq j$.

Thus, without any loss of generality, we will henceforth assume that any H-type group \mathbb{G} is of this form. For an H-type group \mathbb{G} and $x \in \mathbb{G}$, we will therefore use the notation

$$x = (w, z) = (w_1, \dots, w_n, z_1, \dots, z_m),$$

for $w \in \mathbb{R}^n$ and $z \in \mathbb{R}^m$. It is clear that the point $(0, 0)$ is the identity in \mathbb{G} and the inverse operation is $(w, z)^{-1} = (-w, -z)$.

There is a natural family of dilation operators on an H-type group \mathbb{G} :

Definition 2.3. *The family of operators $\delta_\lambda : \mathbb{G} \rightarrow \mathbb{G}$ for $\lambda > 0$ defined by*

$$\delta_\lambda(w, z) = (\lambda w, \lambda z)$$

for $(w, z) \in \mathbb{G}$ is called the family of dilations on \mathbb{G} . The map δ_λ is a group homomorphism, in the sense that $\delta_\lambda(x \circ y) = \delta_\lambda(x) \circ \delta_\lambda(y)$, $\forall x, y \in \mathbb{G}$.

We can identify \mathfrak{g} with the space spanned by the left-invariant vector fields $\{X_1, \dots, X_n, Z_1, \dots, Z_m\}$ on \mathbb{G} , where

$$X_i = \frac{\partial}{\partial w_i} + \frac{1}{2} \sum_{k=1}^m \sum_{l=1}^n U_{il}^{(k)} w_l \frac{\partial}{\partial z_k}, \quad Z_j = \frac{\partial}{\partial z_j}, \quad (2.2)$$

for $i \in \{1, \dots, n\}, j \in \{1, \dots, m\}$. One can calculate that

$$[X_i, X_j] = \sum_{k=1}^m U_{ji}^{(k)} Z_k, \quad (2.3)$$

from which it follows that $\text{span}\{X_i, [X_j, X_k] : i, j, k \in \{1, \dots, n\}\} = \mathfrak{g}$. This is equivalent to saying that the H-type group \mathbb{G} is a Carnot group of step 2.

Definition 2.4. *The second order differential operator $\Delta_{\mathbb{G}} = \sum_{j=1}^n X_j^2$ is the canonical sub-Laplacian on \mathbb{G} . The vector-valued operator $\nabla_{\mathbb{G}} = (X_1, \dots, X_n)$ is the canonical sub-gradient on \mathbb{G} .*

We will treat a general H-type group \mathbb{G} as a metric space, and we can do this in two natural ways. Despite the fact that both metrics are equivalent (in fact all homogeneous metrics on an H-type group are equivalent — see Proposition 5.1.4 of [6]), it turns out that the choice of metric will make a big difference to the spectrum of the associated operator.

The first, and perhaps most common way to define a metric on an H-type group \mathbb{G} is to exploit the structure of the group:

Definition 2.5 (Carnot-Carathéodory distance). *Let $\gamma : [0, 1] \rightarrow \mathbb{G}$ be an absolutely continuous path. We say that γ is horizontal if there exist measurable functions $a_1, \dots, a_n : [0, 1] \rightarrow \mathbb{R}$ such that*

$$\dot{\gamma}(t) = \sum_{i=1}^n a_i(t) X_i(\gamma(t))$$

for almost all $t \in [0, 1]$ i.e. $\dot{\gamma}(t) \in \text{span} \{X_1(\gamma(t)), \dots, X_n(\gamma(t))\}$ almost everywhere. For such a horizontal curve γ , we define the length of γ to be

$$|\gamma| := \int_0^1 \left(\sum_{i=1}^n a_i^2(t) \right)^{\frac{1}{2}} dt.$$

We then define the Carnot-Carathéodory distance $d(x, y)$ between two points $x, y \in \mathbb{G}$ to be

$$d(x, y) := \inf \{ |\gamma| \text{ such that } \gamma : [0, 1] \rightarrow \mathbb{G} \text{ is horizontal and } \gamma(0) = x, \gamma(1) = y \}.$$

We will write $d(x) := d(x, 0)$.

It should be noted that the Carnot-Carathéodory distance is well-defined by Chow's Theorem (see for example [4, 24]).

The second natural distance function that will be of interest to us arises from the fundamental solution of the sub-Laplacian $\Delta_{\mathbb{G}}$. This was discovered by A. Kaplan in [15] on general H-type groups, extending the work of G.B. Folland [11] on the Heisenberg group.

Definition 2.6 (Kaplan distance). *Define the function $N : \mathbb{G} \rightarrow [0, \infty)$ by $N(x) = (|w|^4 + 16|z|^2)^{1/4}$ for $x = (w, z) \in \mathbb{G}$. Then N is a symmetric homogeneous norm, which we will call the Kaplan distance. Moreover, $F = N^{2-Q}$ is a fundamental solution of $\Delta_{\mathbb{G}}$, where $Q = 2m + n$ is the homogeneous dimension of the group, in the sense that F is smooth out of the origin and*

$$\Delta_{\mathbb{G}} F(x) = 0 \quad \text{in} \quad \mathbb{G} \setminus \{0\}.$$

An important difference between the two distance functions is that N is smooth everywhere outside the origin, whilst d has points of non-smoothness on the centre of the group:

Proposition 2.7. (i) $d : \mathbb{G} \rightarrow [0, \infty)$ is smooth on the set $\{x = (w, z) \in \mathbb{G} : w \neq 0\}$, and $|\nabla_{\mathbb{G}} d(x)| = 1$, for all $x = (w, z) \in \mathbb{G}$ such that $w \neq 0$.

(ii) $N : \mathbb{G} \rightarrow [0, \infty)$ is smooth on $\mathbb{G} \setminus \{0\}$, and $|\nabla_{\mathbb{G}} N(x)| = \|x\|/N(x)$, for all $x = (w, z) \in \mathbb{G}$ such that $x \neq 0$, where $\|x\| := |w| = (\sum_{i=1}^n w_i^2)^{\frac{1}{2}}$.

Proof. Part (i) is found in [19], whilst part (ii) can be calculated directly. \square

Example 2.8 (The Heisenberg group). *The main example of an H-type group to keep in mind is the Heisenberg group \mathbb{H} . In fact H-type groups were introduced as a generalisation of the Heisenberg group. \mathbb{H} can be realised as \mathbb{R}^{2+1} with the group operation*

$$(w_1, w_2, z) \circ (\omega_1, \omega_2, \zeta) = \left(w_1 + \omega_1, w_2 + \omega_2, z + \zeta + \frac{1}{2}(w_1\omega_2 - w_2\omega_1) \right)$$

for $w = (w_1, w_2), \omega = (\omega_1, \omega_2) \in \mathbb{R}^2$ and $z, \zeta \in \mathbb{R}$. The left-invariant vector fields on \mathbb{H} are given by

$$X_1 = \partial_{w_1} - \frac{1}{2}w_2\partial_z, \quad X_2 = \partial_{w_2} + \frac{1}{2}w_1\partial_z, \quad Z = \partial_z,$$

and one can easily calculate that $[X_1, X_2] = Z$ and $[X_1, Z] = [X_2, Z] = 0$. Higher dimensional Heisenberg groups can similarly be defined, and all have the common characteristic of a one-dimensional centre.

3 Generalisation of a classical result to the Heisenberg group

We begin our investigation into the spectrum of sub-elliptic operators of the form (1.2) by working in the Heisenberg group, \mathbb{H} . The aim of this section is to prove a direct analogue of the following classical result for Schrödinger operators on \mathbb{R}^n (see for example Theorem XIII.67 of [21]): consider the operator $H = \Delta + V$ on \mathbb{R}^n , where Δ is the standard Laplacian on \mathbb{R}^n and $V \in L^1_{loc}(\mathbb{R}^n)$ is a potential which is bounded from below. Then, if $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$, it follows that H has a purely discrete spectrum.

We will utilise a useful representation of the sub-Laplacian $\Delta_{\mathbb{H}}$ on \mathbb{H} given in [13]. Indeed, denote by \mathcal{F}_3 the partial Fourier transform with respect to the third variable:

$$\mathcal{F}_3 f(w_1, w_2, \zeta) := (2\pi)^{-1/2} \int_{-\infty}^{\infty} e^{-iz\zeta} f(w_1, w_2, z) dz,$$

for $x = (w, z) \in \mathbb{H}$. Then it can be shown that

$$\mathcal{F}_3(-\Delta_{\mathbb{H}} f)(w, \zeta) = (i\nabla_w + \zeta \mathbf{A}(w))^2 \mathcal{F}_3 f(w, \zeta),$$

where $w = (w_1, w_2) \in \mathbb{R}^2$, $\nabla_w = (\partial_{w_1}, \partial_{w_2})$ and $\mathbf{A}(w) = \frac{1}{2}(-w_2, w_1)$. For fixed ζ the operator $(i\nabla_w + \zeta \mathbf{A}(w))^2$ has been well studied, since it corresponds to the Hamiltonian of a particle moving in a magnetic field (see [16]). Indeed, we have the following spectral decomposition:

$$\mathcal{F}_3(-\Delta_{\mathbb{H}} f)(w, \zeta) = \sum_{k=0}^{\infty} \lambda_k(\zeta) \mathcal{P}_k \mathcal{F}_3 f(w, \zeta), \quad w \in \mathbb{R}^2, \zeta \in \mathbb{R}$$

where $\lambda_k(\zeta) := |\zeta|(2k+1)$, $k \in \{0, 1, \dots\}$, and \mathcal{P}_k is the orthogonal eigenprojection given by

$$\mathcal{P}_k f(w) = \int_{\mathbb{R}^2} f(w') \pi_k(w, w') dw',$$

for $w \in \mathbb{R}^2$, where

$$\pi_k(w, w') = \frac{|\zeta|}{2\pi} e^{-\frac{|\zeta|}{2}i(w_1 w'_2 - w_2 w'_1) - \frac{|\zeta|}{4}|w - w'|^2} L_k\left(\frac{|\zeta|}{2}|w - w'|^2\right)$$

and L_k is the k -th Laguerre polynomial. If we then define $\hat{f}(\zeta, k) := \|\mathcal{P}_k \mathcal{F}_3 f(w, \zeta)\|_{L^2(dw)}$ for any $f \in L^2(\mathbb{H})$, one can calculate that

$$\int_{\mathbb{H}} f(x)(-\Delta_{\mathbb{H}} f)(x) dx = \sum_{k=0}^{\infty} \int_{\mathbb{R}} |\zeta|(2k+1) \left| \hat{f}(\zeta, k) \right|^2 d\zeta. \quad (3.1)$$

Moreover, by the spectral theorem we can define a functional calculus for the operator $\Delta_{\mathbb{H}}$. Indeed, for any Borel function $\alpha : [0, \infty) \rightarrow \mathbb{R}$, we define

$$\varphi(-\Delta_{\mathbb{H}}) = \varphi(\lambda_k(\zeta)), \quad (3.2)$$

where the right hand side represents the operator $\mathcal{F}_3^* \sum_k \varphi(\lambda_k(\zeta)) \mathcal{P}_k \mathcal{F}_3$, and we have that

$$\int_{\mathbb{H}} f(x) \varphi(-\Delta_{\mathbb{H}}) f(x) dx = \sum_{k=0}^{\infty} \int_{\mathbb{R}} \varphi(|\zeta|(2k+1)) \left| \hat{f}(\zeta, k) \right|^2 d\zeta. \quad (3.3)$$

Finally, for a self-adjoint operator H on $L^2(\mathbb{H})$, let $\theta_m(H)$, $m \in \mathbb{N} \cup \{0\}$ be as in the Min-Max principle i.e. either $\theta_m(H)$ is the m -th eigenvalue below the bottom of the essential spectrum of H , or $\theta_m(H)$ is the bottom of the essential spectrum and $\theta_m(H) = \theta_{m'}(H)$ for $m' \geq m$ (see for example Theorem XIII.1 of [21]).

Theorem 3.1. *Suppose V is in $L^1_{loc}(\mathbb{H})$ and is bounded from below. Suppose also that for every $L > 0$ there exists $R_L > 0$ such that $V(x) \geq L$ whenever $d(x) \geq R_L$. Then the operator $\mathcal{L} = -\Delta_{\mathbb{H}} + V$ on $L^2(\mathbb{H})$ has empty essential spectrum. In particular it has a purely discrete set of eigenvalues and a complete set of eigenfunctions.*

Proof. Suppose W is a bounded function, supported in a compact set $\Omega \subset \mathbb{R}^3$, so that

$$\sup_{x \in \Omega} |W(x)| \leq M,$$

for some $M \in \mathbb{R}$. For $\varepsilon > 0$ consider the operator

$$W\phi_\varepsilon(-\Delta_{\mathbb{H}}),$$

where $\phi_\varepsilon(t) = (\varepsilon t^2 + t + 1)^{-1}$ for $t \in \mathbb{R}$. Using the above observations, and following [13], we have that

$$\begin{aligned} \text{Tr}(W^2\phi_\varepsilon^2(-\Delta_{\mathbb{H}})) &\leq M^2 \frac{1}{2\pi} \int_{\Omega} \int_{-\infty}^{\infty} \sum_{k=0}^{\infty} \phi_\varepsilon^2(\lambda_k(\zeta)) \pi_k(w, w) d\zeta dw \\ &= M^2 \frac{|\Omega|}{4\pi^2} \sum_{k=0}^{\infty} \int_{-\infty}^{\infty} \phi_\varepsilon^2(\lambda_k(\zeta)) |\zeta| d\zeta \\ &= M^2 \frac{|\Omega|}{2\pi^2} \sum_{k=0}^{\infty} \int_0^{\infty} \frac{|\zeta|}{(\varepsilon|\zeta|^2(2k+1)^2 + |\zeta|(2k+1) + 1)^2} d\zeta < \infty. \end{aligned}$$

Since $W\phi_\varepsilon(-\Delta_{\mathbb{H}})$ is positive and self-adjoint on $L^2(\mathbb{H})$, we thus have that $W\phi_\varepsilon(-\Delta_{\mathbb{H}})$ is Hilbert-Schmidt for all $\varepsilon > 0$. Moreover,

$$(\varepsilon\lambda_k(\zeta)^2 + \lambda_k(\zeta) + 1)^{-1} \rightarrow (\lambda_k(\zeta) + 1)^{-1} \equiv \phi_0(\lambda_k(\zeta))$$

in $L^\infty(\mathbb{R}) \times l^\infty(\mathbb{N} \cup \{0\})$ as $\varepsilon \rightarrow 0$. Therefore $W\phi_0(-\Delta_{\mathbb{H}})$ is a norm-limit of Hilbert-Schmidt operators:

$$\begin{aligned} \|W(\phi_0 - \phi_\varepsilon)(-\Delta_{\mathbb{H}})\psi\|_2^2 &\leq M^2 \sum_{k=0}^{\infty} \int_{\mathbb{R}} (\phi_0 - \phi_\varepsilon)^2(\lambda_k(\zeta)) \left| \hat{\psi}(\zeta, k) \right|^2 d\zeta \\ &\leq \varepsilon^2 M^2 \sum_{k=0}^{\infty} \int_{\mathbb{R}} \left| \hat{\psi}(\zeta, k) \right|^2 d\zeta \\ &= \varepsilon^2 M^2 \|\psi\|_2^2. \end{aligned}$$

We can thus conclude that $W\phi_0(-\Delta_{\mathbb{H}})$ is a compact operator, or in other words that W is relatively compact with respect to $-\Delta_{\mathbb{H}}$.

Since $W\phi_0(-\Delta_{\mathbb{H}})$ is compact, by Weyl's Theorem (see Corollary 2 of Theorem XIII.14 of [21]),

$$\sigma_{ess}(-\Delta_{\mathbb{H}} + W) = \sigma_{ess}(-\Delta_{\mathbb{H}}) = [0, \infty).$$

Therefore by the Min-Max principle $\theta_m(-\Delta_{\mathbb{H}} + W) \geq -1$ for m sufficiently large.

Now, given $a > 0$, define V_a by

$$V_a(x) = \min\{V(x), a + 1\} - a - 1.$$

Then V_a has compact support, since $V(x) \rightarrow \infty$ as $d(x) \rightarrow \infty$. Thus, by the above considerations, $\theta_m(-\Delta_{\mathbb{H}} + V_a) \geq -1$ for large m . Finally, since

$$\theta_m(\mathcal{L}) \geq \theta_m(-\Delta_{\mathbb{H}} + V_a) + a + 1,$$

we see that $\theta_m(\mathcal{L}) \geq a$ for large m . Since a is arbitrary we reach the desired conclusion. \square

As mentioned in the introduction, we are actually interested in probability measures of the form $\mu_U(dx) := Z^{-1}e^{-U(x)}dx$ on \mathbb{H} , where $Z = \int_{\mathbb{H}} e^{-U(x)}dx < \infty$, with which we can associate a positive and self-adjoint operator $\mathcal{L} = -\Delta_{\mathbb{H}} + \nabla_{\mathbb{H}}U \cdot \nabla_{\mathbb{H}}$ on $L^2(d\mu_U)$.

Corollary 3.2. *Suppose that $V = \frac{1}{4}|\nabla_{\mathbb{H}}U|^2 - \frac{1}{2}\Delta_{\mathbb{H}}U$ is in $L^1_{loc}(\mathbb{H})$, is bounded from below, and is such that $V(x) \rightarrow \infty$ as $d(x) \rightarrow \infty$. Let $\mathcal{L} = -\Delta_{\mathbb{H}} + \nabla_{\mathbb{H}}U \cdot \nabla_{\mathbb{H}}$, so that \mathcal{L} is a positive self-adjoint operator on $L^2(\mu_U)$. Then $\sigma_{ess}(\mathcal{L}) = \emptyset$.*

Proof. Follows from Theorem 3.1 and the identity

$$\int_{\mathbb{H}} f(-\Delta_{\mathbb{H}} + \nabla_{\mathbb{H}}U \cdot \nabla_{\mathbb{H}})f d\mu = \int_{\mathbb{H}} g(-\Delta_{\mathbb{H}} + V)g dx, \quad g = fe^{-\frac{1}{2}U}.$$

\square

Following a recent trend (see [14] and references therein), for the rest of this paper we will concentrate on the case when U is given as a power of one of the natural distance functions i.e. $U(x) = -\alpha d^p(x)$ or $U(x) = -\alpha N^p(x)$ with $p \in (1, \infty)$ and $\alpha > 0$, where d and N are the Carnot-Carathéodory and Kaplan distance functions respectively. We can therefore try and apply Corollary 3.2 in both these cases. However, this is not straightforward in either case. Indeed, when $U = \alpha d^p$ we have that

$$\frac{1}{4}|\nabla_{\mathbb{H}}U|^2 - \frac{1}{2}\Delta_{\mathbb{H}}U = \frac{\alpha^2 p^2}{4}d^{2(p-1)} - \frac{\alpha p(p-1)}{2}d^{p-2} - \frac{\alpha p}{2}d^{p-1}\Delta_{\mathbb{H}}d, \quad (3.4)$$

but we must understand this expression in the sense of distributions since $\Delta_{\mathbb{H}}d$ is not defined on the centre of the group. Moreover, when $U = \alpha N^p$ we have

$$\begin{aligned} \frac{1}{4}|\nabla_{\mathbb{H}}U|^2 - \frac{1}{2}\Delta_{\mathbb{H}}U &= \frac{\alpha^2 p^2}{4}N^{2p-4}(x)\|x\|^2 \\ &\quad - \left(\frac{\alpha p(p-1)}{2} + \frac{3\alpha p}{2}\right)N^{p-4}(x)\|x\|^2, \end{aligned}$$

where $\|x\|^2 = w_1^2 + w_2^2$ as usual, so that it is certainly not true that $V(x) \rightarrow \infty$ as $|x| \rightarrow \infty$ ($V(x) = 0$ for all $x = (0, z) \in \mathbb{H}$). We therefore look for alternative methods to investigate these interesting cases.

4 Spectral information via functional inequalities

The relationship between functional inequalities and the spectrum of operators is a very interesting and much studied one. Indeed, if (Ω, μ) is a probability space and $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ is a positive self-adjoint operator on $L^2(\mu)$, then it is well-known that \mathcal{L} has a gap at the bottom of its spectrum if and only if there exists a constant $c_0 > 0$ such that

$$\mu(f - \mu(f))^2 \leq c_0 \mathcal{E}(f, f),$$

where $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ is the Dirichlet form associated to \mathcal{L} . More recently this relationship has been further illustrated by the work of F. Cipriani ([7]) and F. Y. Wang ([25]) in which functional inequalities are introduced that characterise the essential spectra of operators under very general conditions. In this section we aim to use functional inequalities to overcome the problems encountered at the end of section 3 in the more general setting of H-type groups.

4.1 Super-Poincaré inequalities

To state the results of Wang and Cipriani in full generality, we first need the following two technical definitions.

Definition 4.1. *A topological space Ω is a Lusin space if Ω is homeomorphic to a Borel subset of a compact metric space.*

Remark 4.2. *It should be noted that, as shown in Theorem 82.5 of [22], every complete metric space is a Lusin space. In particular, any H-type group \mathbb{G} is a Lusin space.*

Definition 4.3. Let Ω be a Lusin space, and μ a positive Radon measure on Ω having full topological support. A positive, self-adjoint operator $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ on $L^2(\mu)$, with associated closed Dirichlet form $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ defined by

$$\mathcal{E}(f, g) = \mu(f\mathcal{L}g), \quad f, g \in \mathcal{D}(\mathcal{L}),$$

is called a Persson operator if $\inf \sigma_{ess}(\mathcal{L}) = \sup \{\Sigma(K) : K \subset \Omega \text{ is compact}\}$, where

$$\Sigma(K) := \inf \left\{ \frac{\mathcal{E}(f, f)}{\|f\|_2^2} : f \in \mathcal{D}(\mathcal{E}), \text{ supp}(f) \subset K^c \right\}.$$

This class of operator was introduced by A. Persson in [20]. The result below is a combination of the independent work of Wang and Cipriani, and is explicitly stated in [26].

Theorem 4.4 (Wang/Cipriani). Let Ω be a Lusin space, μ a positive Radon measure on Ω having full topological support, and $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ a Persson operator on $L^2(\mu)$. Then the inequality

$$\mu(f^2) \leq r\mu(f\mathcal{L}f) + \beta(r)(\mu|f|)^2, \quad \forall r > r_0, \quad f \in \mathcal{D}(\mathcal{L}), \quad (4.1)$$

for some decreasing function $\beta : (r_0, \infty) \rightarrow (0, \infty)$ and $r_0 \geq 0$ holds if and only if $\sigma_{ess}(\mathcal{L}) \subset [r_0^{-1}, \infty)$. In particular, (4.1) is satisfied with $r_0 = 0$ if and only if $\sigma_{ess}(\mathcal{L}) = \emptyset$.

Inequality (4.1) is known as a *super-Poincaré inequality*. In a similar way to the generalisation of the standard logarithmic Sobolev inequality to the LS_q inequality (see [5]), we can generalise the super-Poincaré inequality to a *q-super-Poincaré inequality*:

Definition 4.5. Let (Ω, μ) be a probability space, equipped with a metric $d : \Omega \times \Omega \rightarrow [0, \infty)$. For $q \in (1, 2]$, we say that μ satisfies a *q-super-Poincaré inequality*, or SP_q for short, with constant r_0 , if

$$\mu|f|^q \leq r\mu|\nabla f|^q + \beta(r) \left(\mu|f|^{\frac{q}{2}} \right)^2, \quad \forall r > r_0, \quad (SP_q) \quad (4.2)$$

for all locally Lipschitz functions f and some $\beta : (r_0, \infty) \rightarrow (0, \infty)$, where $|\nabla f|(x) \equiv \limsup_{d(x,y) \rightarrow 0} |f(x) - f(y)|/d(x, y)$.

Remark 4.6. In an H -type group \mathbb{G} equipped with the Carnot-Carathéodory distance d , $|\nabla_{\mathbb{G}} f|(x) = \limsup_{d(x,y) \rightarrow 0} |f(x) - f(y)|/d(x, y)$.

4.2 Potentials defined by the Carnot-Carathéodory distance

Let \mathbb{G} be an H-type group as usual, equipped with the Carnot-Carathéodory distance d . Let μ_p be the probability measure on \mathbb{G} defined by

$$\mu_p(dx) := \frac{e^{-\alpha d^p(x)}}{Z} dx \quad (4.3)$$

where $Z = \int e^{-\alpha d^p(x)} dx$ is the normalisation constant, and $p \in (1, \infty), \alpha > 0$. Define

$$\mathcal{L}_p := -\Delta_{\mathbb{G}} + \nabla_{\mathbb{G}}(\alpha d^p) \cdot \nabla_{\mathbb{G}} = -\Delta_{\mathbb{G}} + \alpha p d^{p-1} \nabla_{\mathbb{G}} d \cdot \nabla_{\mathbb{G}} \quad (4.4)$$

as a positive self-adjoint operator acting on $L^2(\mu_p)$. The associated Dirichlet form $\mathcal{E}_p(f, g)$ is then given by

$$\mathcal{E}_p(f, g) = \mu_p(f \mathcal{L}_p g) = \int_{\mathbb{G}} \nabla_{\mathbb{G}} f \cdot \nabla_{\mathbb{G}} g d\mu_p. \quad (4.5)$$

We aim to prove the following:

Theorem 4.7. *For any $p > 1$ the positive self-adjoint operator \mathcal{L}_p on $L^2(\mu_p)$ given by (4.4) has a purely discrete spectrum i.e. $\sigma_{ess}(\mathcal{L}_p) = \emptyset$.*

The idea is to use Theorem 4.4. It is clear that we first need to show that \mathcal{L}_p is a Persson operator. We make use of the following very general result stated in the setting of Dirichlet forms, proved by G. Grillo in [12] (and also stated explicitly in [7]).

Theorem 4.8 (Grillo). *Let (Ω, μ) be a locally compact, separable metric space, and $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ a regular, strongly local Dirichlet form on $L^2(\Omega)$, with associated positive self-adjoint operator $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$.*

Define the associated intrinsic pseudo-metric ρ on Ω by

$$\rho(x, y) := \sup \{ |f(x) - f(y)| : f \in \mathcal{D}(\mathcal{E}) \cap C_0(\Omega), \Gamma(f, f) \leq 1 \}$$

where for $f \in \mathcal{D}(\mathcal{E})$, $\Gamma(f, f)$ is such that

$$\int_{\Omega} g \Gamma(f, f) d\mu = \mathcal{E}(gf, f) - \frac{1}{2} \mathcal{E}(f^2, g), \quad f, g \in \mathcal{D}(\mathcal{E}) \cap C_0(\Omega).$$

Suppose ρ is a true metric generating the original topology of Ω . Then the operator $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ is a Persson operator.

Corollary 4.9. *The operator \mathcal{L}_p given by (4.4) acting on $L^2(\mu_p)$ is a Persson operator.*

Proof. The intrinsic pseudo-metric associated to \mathcal{L}_p is given by

$$\rho(x, y) = \sup \{ |f(x) - f(y)| : |\nabla_{\mathbb{G}} f|^2 \leq 1 \},$$

which is nothing more than the Carnot-Carathéodory distance (by definition). \square

The next result we prove on route to Theorem 4.7 is that the measures μ_p satisfy certain super-Poincaré inequalities.

Theorem 4.10. *Let μ_p be the probability measure on \mathbb{G} given by (4.3).*

(i) *Suppose $p \geq 2$. Then μ_p satisfies an SP_q inequality with constant $r_0 = 0$ i.e.*

$$\mu_p |f|^q \leq r \mu_p |\nabla_{\mathbb{G}} f|^q + \beta(r) \left(\mu_p |f|^{\frac{q}{2}} \right)^2, \quad \forall r > 0,$$

where $\frac{1}{p} + \frac{1}{q} = 1$, for some function $\beta : (0, \infty) \rightarrow (0, \infty)$ and for all locally Lipschitz functions f .

(ii) *Suppose $p \in (1, 2]$. Then μ_p satisfies an SP_2 inequality with constant $r_0 = 0$ i.e.*

$$\mu_p(f^2) \leq r \mu_p |\nabla_{\mathbb{G}} f|^2 + \beta(r) (\mu_p |f|)^2, \quad \forall r > 0,$$

for some function $\beta : (0, \infty) \rightarrow (0, \infty)$ and for all locally Lipschitz functions f .

Proof. The idea is to pass from a logarithmic Sobolev inequality for the measure μ_p , which is true by Theorem 4.1 of [14], to a super-Poincaré inequality by adapting the methods of [25].

We first deal with the case $p \geq 2$. Without loss of generality we may assume that $f \geq 0$. By Theorem 4.1 of [14], we have that there exists a constant c such that

$$\mu_p \left(f^q \log \frac{f^q}{\mu_p f^q} \right) \leq c \mu_p |\nabla_{\mathbb{G}} f|^q \quad (4.6)$$

where $\frac{1}{p} + \frac{1}{q} = 1$. Let $g : (0, \infty) \rightarrow \mathbb{R}$ be given by $g(\xi) = t\xi - \xi \log \left(\frac{\xi^2}{a} \right)$ for any $t, a > 0$. We then have that

$$\max_{\{\xi > 0\}} g(\xi) = 2\sqrt{ae^{t-2}}. \quad (4.7)$$

Suppose that $\mu_p(f^{\frac{q}{2}}) = 1$, and set $a = \mu_p(f^q)$. Then by (4.7), for all $t > 0$,

$$\begin{aligned} t f^{\frac{q}{2}} - f^{\frac{q}{2}} \log \left(\frac{f^q}{a} \right) &\leq 2\sqrt{ae^{t-2}} \\ \Rightarrow \mu_p \left(f^q \log \frac{f^q}{a} \right) &\geq ta - 2\sqrt{ae^{t-2}}, \end{aligned} \quad (4.8)$$

using the fact that $f \geq 0$ and $\mu_p(f^{\frac{q}{2}}) = 1$. Setting $b = \mu_p|\nabla_{\mathbb{G}}f|^q$, by (4.6), we then have

$$ta - 2\sqrt{ae^{t-2}} - cb \leq 0.$$

Solving this quadratic inequality gives

$$\sqrt{a} \leq \frac{2\sqrt{e^{t-2}}}{2t} + \frac{\sqrt{4e^{t-2} + 4tcb}}{2t}$$

for $t > 0$, so that $a \leq \frac{2c}{t}b + 4\frac{e^{t-2}}{t^2}$. Thus

$$\mu_p(f^q) \leq \frac{2c}{t}\mu_p|\nabla_{\mathbb{G}}f|^q + 4\frac{e^{t-2}}{t^2},$$

for all $t > 0$ and f such that $\mu_p(f^{\frac{q}{2}}) = 1$. Replacing f by $\frac{f}{\mu_p(f^{q/2})^{2/q}}$ yields

$$\mu_p(f^q) \leq \frac{2c}{t}\mu_p|\nabla_{\mathbb{G}}f|^q + 4\frac{e^{t-2}}{t^2} \left(\mu_p(f^{\frac{q}{2}})\right)^2$$

for all $t > 0$. Taking $r = \frac{2c}{t}$ we see that SP_q holds, proving part (i).

In the case where $p \in (1, 2)$, we no longer have an inequality of the type (4.6). However, by Theorem 4.3 of [14] there exists a constant $c \in (0, \infty)$ such that

$$\mu_p \left(f^2 [\log(1 + f^2)]^\theta \right) \leq c\mu_p|\nabla_{\mathbb{G}}f|^2 + (\log 2)^\theta, \quad \mu_p(f^2) = 1, \quad (4.9)$$

where $\theta = \frac{2(p-1)}{p}$. In this case we instead let $g : (0, \infty) \rightarrow \mathbb{R}$ be given by $g(\xi) = t\xi - \xi \left[\log \left(1 + \frac{\xi^2}{a} \right) \right]^\theta$ for $t, a > 0$, so that $\sup_{\{\xi > 0\}} g(\xi) \leq t\sqrt{a(e^{t^{1/\theta}} - 1)}$. Proceeding now in a very similar way as in the proof of part (i), we arrive at an SP_2 inequality. \square

The final result we need is that SP_q inequalities are stronger than SP_2 inequalities (at least when the dimension of the underlying space is finite).

Lemma 4.11. *Suppose an arbitrary probability measure μ on \mathbb{G} satisfies an SP_q inequality with $q \in (1, 2]$ and constant $r_0 = 0$. Then μ also satisfies an SP_2 inequality with constant $r_0 = 0$.*

Proof. As usual, without loss of generality we may suppose $f \geq 0$. Let $q < 2$ (there is nothing to prove if $q = 2$). Applying the SP_q inequality to $f^{\frac{2}{q}}$ yields,

$$\mu(f^2) \leq r\mu \left| \nabla_{\mathbb{G}}f^{\frac{2}{q}} \right|^q + \beta(r)(\mu f)^2, \quad \forall r > 0.$$

Therefore for all $r > 0$, we have by Hölder's inequality followed by Young's inequality,

$$\begin{aligned}\mu(f^2) &\leq \frac{2^q r}{q^q} \mu(f^{2-q} |\nabla_{\mathbb{G}} f|^q) + \beta(r) (\mu f)^2 \\ &\leq \frac{2^{q-1} r}{q^{q-1}} \tau^{\frac{2-q}{q}} \mu |\nabla_{\mathbb{G}} f|^2 + \frac{2^{q-1} r (2-q)}{q^q} \tau^{-1} \mu(f^2) + \beta(r) (\mu f)^2\end{aligned}$$

for all $\tau > 0$. Taking $\tau = \frac{2^q r (2-q)}{q^q}$ we see that

$$\begin{aligned}\frac{1}{2} \mu(f^2) &\leq \frac{2^{q-1} r}{q^{q-1}} \tau^{\frac{2-q}{q}} \mu |\nabla_{\mathbb{G}} f|^2 + \beta(r) (\mu f)^2 \\ &= \frac{2^{q-1}}{q^{q-1}} \left(\frac{2}{q} \right)^{2-q} (2-q)^{\frac{2-q}{q}} r^{\frac{2}{q}} \mu |\nabla_{\mathbb{G}} f|^2 + \beta(r) (\mu f)^2 \\ \Rightarrow \mu(f^2) &\leq \frac{4}{q} (2-q)^{\frac{2-q}{q}} r^{\frac{2}{q}} \mu |\nabla_{\mathbb{G}} f|^2 + 2\beta(r) (\mu f)^2.\end{aligned}$$

Taking $s = \frac{4}{q} (2-q)^{\frac{2-q}{q}} r^{\frac{2}{q}}$ then yields the result. \square

Proof of Theorem 4.7. We can now combine all of the above results to arrive at Theorem 4.7. Indeed by Theorem 4.10 and Lemma 4.11, we have that the measures μ_p satisfy a super-Poincaré inequality with constant $r_0 = 0$ for all $p > 1$. Moreover, by Corollary 4.9, \mathcal{L}_p is a Persson operator, so that we may conclude by applying Theorem 4.4. \square

Remark 4.12. *Since the main result we use to prove Theorem 4.7 is that the measure μ_p satisfies a logarithmic Sobolev inequality, we can greatly extend the class of measures for which these results remain valid when $p \geq 2$. Indeed, if we define*

$$\tilde{\mu}_p = \frac{e^{-W-V}}{Z'} d\mu_p$$

with W a differentiable potential satisfying $|\nabla_{\mathbb{G}} W|^q \leq d^\gamma + K$ for constants K and $\gamma \in (0, p)$, and V a measurable function such that $\max V - \min V < \infty$, then all the above results remain true for $\tilde{\mu}_p$. This follows from Corollary 4.1 of [14].

Corollary 4.13. *Let $\rho_t(x, y)$ be the heat kernel at time t on an H -type group \mathbb{G} i.e. $\rho_t(x, y)$ is the function (smooth by Hörmander's theorem) such that*

$$e^{t\Delta_{\mathbb{G}}} f(x) = \int_{\mathbb{G}} \rho_t(x, y) f(y) dy.$$

Let $\rho(x) := \rho_1(x, e)$ and define

$$\mathcal{L}_H := -\Delta_{\mathbb{G}} + \nabla_{\mathbb{G}} \log \rho \cdot \nabla_{\mathbb{G}}.$$

Then \mathcal{L}_H is a positive self-adjoint operator on $L^2(\mu_H)$, where $\mu_H(dx) := \rho(x)dx$, and $\sigma_{\text{ess}}(\mathcal{L}_H) = \emptyset$, so that \mathcal{L}_H has a purely discrete spectrum.

Remark 4.14. \mathcal{L}_H can be regarded as the natural Ornstein-Uhlenbeck generator on \mathbb{G} , as suggested in [3], and the resulting Markov process is the natural OU-process associated to the hypoelliptic diffusion on \mathbb{G} .

Proof. It follows exactly as above, once we have recalled that μ_H satisfies a logarithmic Sobolev inequality (see [1, 14] and [17]). \square

4.3 Potentials defined by the Kaplan distance

An interesting question to ask is whether one can replace the Carnot-Carathéodory distance with the Kaplan distance in the above work. At first glance such a question might seem simple, since all homogeneous metrics on \mathbb{G} are equivalent. However, as we will see, this is not the case, and there are some fundamental differences between the two situations which arise from the different smoothness properties of the two distance functions.

To proceed, suppose now that we are working in an H-type group \mathbb{G} as before, but now equipped with a probability measure and associated self-adjoint operator given by

$$\nu_p(dx) := \frac{e^{-\alpha N^p(x)}}{Z} dx, \quad \mathcal{T}_p = -\Delta_{\mathbb{G}} + \alpha p N^{p-1} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} \quad (4.10)$$

respectively, where $p \in (1, \infty)$, $\alpha > 0$ and $Z = \int e^{-\alpha N^p(x)} dx$.

Trying to apply the functional inequality approach of section 4.1, we immediately come up against a problem in the form of Theorem 6.3 from [14]:

Theorem 4.15 (Hebisch-Zegarliński). *The measure ν_p on \mathbb{G} given by (4.10) with $p > 1$ satisfies no LS_q inequality with $q \in (1, 2]$.*

Thus we cannot simply follow the proof of Theorem 4.7 to conclude that the operator \mathcal{T}_p given by (4.10) has empty essential spectrum. Theorem 4.15 illustrates a major difference in the behaviour of the measures defined with the Carnot-Carathéodory distance and those defined with the Kaplan distance.

In view of this, it seems that the problem of gaining spectral information about the operator \mathcal{T}_p given by (4.10) is an interesting one. We therefore start by asking whether such operators have a spectral gap. This question is completely answered by Theorems 4.16 and 4.19 below.

Theorem 4.16. *If $p < 2$, then the measure ν_p given by (4.10) does not satisfy a spectral gap inequality. In particular the operator \mathcal{T}_p given by (4.10) does not have a spectral gap, and hence it does not have empty essential spectrum.*

To prove this, we make use of the following result, quoted from [14] (Lemma 6.3), and we refer the reader to this work for the proof.

Lemma 4.17. *Let f be a smooth function on \mathbb{G} and d the Carnot-Carathéodory distance as usual. Then at points $x_0 \in \mathbb{G}$ such that $(\nabla_{\mathbb{G}} f)(x_0) = 0$ we have*

$$|f(x) - f(x_0)| \leq O(d^2(x, x_0))$$

for all $x \in \mathbb{G}$.

Proof of Theorem 4.16. Let $p < 2$ and suppose for a contradiction that there exists a constant c_0 such that

$$\nu_p(f^2) - (\nu_p f)^2 \leq c_0 \nu_p |\nabla_{\mathbb{G}} f|^2 \quad (4.11)$$

for all locally Lipschitz functions f .

Fix $x_0 = (0, z) \in \mathbb{G}$ for $z \in \mathbb{R}^m \setminus \{0\}$. Then $|\nabla_{\mathbb{G}} N(x_0)| = \frac{\|x_0\|}{N(x_0)} = 0$ by Proposition 2.7, so that $\nabla_{\mathbb{G}} N(x_0) = 0$. Similarly $\nabla_{\mathbb{G}} N(-x_0) = 0$. Let $r_0 > 0$ be small enough so that $0 \notin B_{r_0}(x_0) = \{y' \in \mathbb{G} : d(y', x_0) \leq r_0\}$. Then N is smooth on $B_{r_0}(x_0)$, and by Lemma 4.17 there exists a constant C_1 such that

$$|N(y) - N(x_0)| \leq C_1 r_0^2, \quad (4.12)$$

for all $y \in B_{r_0}(x_0)$. The same holds for $y \in B_{r_0}(-x_0)$. We now dilate by a factor of $t > 0$. Since N is homogeneous, we have that

$$|N(y) - N(\delta_t(x_0))| = t |N(\delta_{t^{-1}}(y)) - N(x_0)| \leq C_1 t r_0^2$$

for $\delta_{t^{-1}}(y) \in B_{r_0}(x_0) \Leftrightarrow y \in B_{tr_0}(\delta_t(x_0))$, where $(\delta_t)_{t>0}$ is the natural family of dilations given by Definition 2.3. The same holds for $y \in B_{tr_0}(\delta_t(-x_0))$.

Let $r = tr_0$. We have for $y \in B_r(\delta_t(x_0))$ or $y \in B_r(\delta_t(-x_0))$

$$\begin{aligned} |N^p(y) - N^p(\delta_t(x_0))| &\leq C_2 N^{p-1}(\delta_t(x_0)) |N(y) - N(\delta_t(x_0))| \\ &\leq C_3 t^{p-1} t r_0^2 = C_3 t^p r_0^2 \end{aligned}$$

for some constants C_2, C_3 , using the mean value theorem. Thus if we take t large enough so that $r_0 = t^{-\frac{p}{2}}$, we have

$$|N^p(y) - N^p(\delta_t(x_0))| \leq C_3, \quad \forall y \in B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0)),$$

so that

$$\left| \frac{e^{-\beta N^p(y)}}{e^{-\beta N^p(\delta_t(x_0))}} \right| \approx 1 \quad (4.13)$$

for all $y \in B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0))$. Now define

$$\begin{aligned} \varphi(y) = & \max \left\{ \min \left\{ 2 - \frac{N(y, \delta_t(x_0))}{r}, 1 \right\}, 0 \right\} \\ & - \max \left\{ \min \left\{ 2 - \frac{N(y, \delta_t(-x_0))}{r}, 1 \right\}, 0 \right\}. \end{aligned} \quad (4.14)$$

Then φ is a Lipschitz function supported on balls of radius r centred at $\delta_t(x_0)$ and $\delta_t(-x_0)$, which is equal to 1 on balls of radius $r/2$ around these two points and decays to zero linearly in between $r/2$ and r . We can note that by construction, and since the measure ν_p is symmetric about the origin,

$$\int_{\mathbb{G}} \varphi(y) d\nu_p(y) = 0.$$

Applying the spectral gap inequality (4.11) to the function φ , then yields

$$\int_{B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0))} \varphi^2(y) d\nu_p(y) \leq c_0 \int_{B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0))} |\nabla_{\mathbb{G}} \varphi(y)|^2 d\nu_p(y). \quad (4.15)$$

Now, using (4.13), there exist positive constants C_4 and C_5 such that

$$\int_{B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0))} \varphi^2(y) d\nu_p(y) \geq 2 \int_{B_{\frac{r}{2}}(\delta_t(x_0))} d\nu_p(y) \geq C_4 r^Q e^{-\beta t^p N^p(x_0)},$$

and

$$\int_{B_r(\delta_t(x_0)) \cup B_r(\delta_t(-x_0))} |\nabla_{\mathbb{G}} \varphi(y)|^2 d\nu_p(y) \leq 2r^{-2} \int_{B_r(\delta_t(x_0))} d\nu_p(y) \leq C_5 r^{-2+Q} e^{-\beta t^p N^p(x_0)},$$

where $Q = n + 2m$ is the homogeneous dimension of the group. Using these estimates in (4.15) yields

$$C_4 \leq c_0 C_5 r^{-2},$$

where $r = tr_0 = t^{1-\frac{p}{2}}$. Since $p < 2$ and t can be taken arbitrarily large, this is a contradiction. \square

Remark 4.18. *Theorem 4.16 provides another illustration of a fundamental difference between the operators \mathcal{L}_p defined by (4.4) with the Carnot-Carathéodory distance and the operators \mathcal{T}_p defined by (4.10) with the Kaplan distance. Indeed, with $p \in (1, 2)$, by Theorem 4.7, \mathcal{L}_p has empty essential spectrum, while \mathcal{T}_p does not even have a spectral gap.*

Theorem 4.19. *If $p \geq 2$, the measure ν_p given by (4.10) satisfies a q -spectral gap inequality, i.e. there exists a constant c_0 such that*

$$\nu_p |f - \nu_p f|^q \leq c_0 \nu_p |\nabla_{\mathbb{G}} f|^q$$

for all locally Lipschitz functions f , where $\frac{1}{q} + \frac{1}{p} = 1$. In particular, for $p \geq 2$ the operator \mathcal{T}_p associated to ν_p given by (4.10) has a spectral gap.

To prove this we adapt the methods of Hebisch and Zegarliński in [14], and proceed through an intermediate inequality which is similar to the U -bound studied there.

Lemma 4.20. *For $p \geq 2$ there exist constants A, B such that*

$$\nu_p(f^q N^{p-2} \|\cdot\|^2) \leq A \nu_p |\nabla_{\mathbb{G}} f|^q + B \nu_p |f|^q \quad (4.16)$$

for all locally Lipschitz functions f , where $\frac{1}{p} + \frac{1}{q} = 1$, and $\|x\| = |w|$ for $x = (w, z) \in \mathbb{G} = \mathbb{R}^n \times \mathbb{R}^m$.

Proof. We can suppose as usual that $f \geq 0$ and moreover that $f \in C_0^\infty(\mathbb{G})$. We can write

$$(\nabla_{\mathbb{G}} f) e^{-\alpha N^p} = \nabla_{\mathbb{G}} (f e^{-\alpha N^p}) + \alpha p f N^{p-1} (\nabla_{\mathbb{G}} N) e^{-\alpha N^p}.$$

We now take the inner product of both sides of this equation with $\frac{N}{\|x\|} \nabla_{\mathbb{G}} N$ and integrate over \mathbb{G} to arrive at

$$\begin{aligned} \int \frac{N}{\|x\|} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} f e^{-\alpha N^p} dx &= \int \frac{N}{\|x\|} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} (f e^{-\alpha N^p}) dx \\ &\quad + \alpha p \int f \frac{N^p}{\|x\|} |\nabla_{\mathbb{G}} N|^2 e^{-\alpha N^p} dx. \end{aligned}$$

By the Cauchy-Schwarz inequality, we then have

$$\begin{aligned} &\int \frac{N}{\|x\|} |\nabla_{\mathbb{G}} N| |\nabla_{\mathbb{G}} f| e^{-\alpha N^p} dx \\ &\geq \int \frac{N}{\|x\|} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} (f e^{-\alpha N^p}) dx + \alpha p \int f \frac{N^p}{\|x\|} |\nabla_{\mathbb{G}} N|^2 e^{-\alpha N^p} dx, \end{aligned}$$

so that by Proposition 2.7 and integration by parts,

$$\begin{aligned} \int |\nabla_{\mathbb{G}} f| e^{-\alpha N^p} dx &\geq \int \frac{N}{\|x\|} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} (f e^{-\alpha N^p}) dx + \alpha p \int f N^{p-2} \|x\| e^{-\alpha N^p} dx \\ &= - \int f \nabla_{\mathbb{G}} \cdot \left(\frac{N}{\|x\|} \nabla_{\mathbb{G}} N \right) e^{-\alpha N^p} dx + \alpha p \int f N^{p-2} \|x\| e^{-\alpha N^p} dx. \end{aligned} \quad (4.17)$$

Note that

$$\begin{aligned}\nabla_{\mathbb{G}} \cdot \left(\frac{N}{\|x\|} \nabla_{\mathbb{G}} N \right) &= \frac{|\nabla_{\mathbb{G}} N|^2}{\|x\|} + \frac{N}{\|x\|} \Delta_{\mathbb{G}} N - \frac{N}{\|x\|^2} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} \|x\| \\ &= \frac{\|x\|}{N^2} + (Q-1) \frac{\|x\|}{N^2} - \frac{N}{\|x\|^2} \nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} \|x\|. \end{aligned} \quad (4.18)$$

Moreover, denoting $x = (w, z) \in \mathbb{G}$ and recalling the results from Section 2, it can be calculated that

$$\nabla_{\mathbb{G}} N \cdot \nabla_{\mathbb{G}} \|x\| = \frac{\|x\|^3}{N^3}.$$

Using this in (4.18) yields

$$\nabla_{\mathbb{G}} \cdot \left(\frac{N}{\|x\|} \nabla_{\mathbb{G}} N \right) = (Q-1) \frac{\|x\|}{N^2}. \quad (4.19)$$

Putting (4.19) in (4.17) then gives

$$\alpha p \nu_p(f N^{p-2} \|\cdot\|) \leq \nu_p |\nabla_{\mathbb{G}} f| + (Q-1) \nu_p \left(f \frac{\|\cdot\|}{N^2} \right).$$

Replacing f by $f \|\cdot\|$, we see that

$$\alpha p \nu_p(f N^{p-2} \|\cdot\|^2) \leq \nu_p (\|\cdot\| |\nabla_{\mathbb{G}} f|) + Q \nu_p(f), \quad (4.20)$$

using the fact that $\|x\| \leq N(x)$ and $|\nabla_{\mathbb{G}} \|x\|| = 1$. Now, by replacing f by f^q with $\frac{1}{q} + \frac{1}{p} = 1$ in (4.20), we then arrive at

$$\begin{aligned}\alpha p \nu_p(f^q N^{p-2} \|\cdot\|^2) &\leq q \nu_p (\|\cdot\| f^{q-1} |\nabla_{\mathbb{G}} f|) + Q \nu_p(f^q) \\ &\leq \frac{1}{\varepsilon^{q-1}} \nu_p |\nabla_{\mathbb{G}} f|^q + \frac{q}{p} \varepsilon \nu_p (\|\cdot\|^p f^q) + Q \nu_p(f^q),\end{aligned}$$

for all $\varepsilon > 0$, using Young's inequality. Thus

$$\alpha p \nu_p(f^q N^{p-2} \|\cdot\|^2) \leq \frac{1}{\varepsilon^{q-1}} \nu_p |\nabla_{\mathbb{G}} f|^q + \frac{q}{p} \varepsilon \nu_p (N^{p-2} \|\cdot\|^2 f^q) + Q \nu_p(f^q),$$

so that, by taking $\varepsilon < \frac{p^2}{q} \alpha$, we see that

$$\nu_p(f^q N^{p-2} \|\cdot\|^2) \leq A \nu_p |\nabla_{\mathbb{G}} f|^q + B \nu_p(f^q),$$

with $A = \varepsilon^{-q+1}(\alpha p - \frac{q}{p} \varepsilon)^{-1}$, and $B = Q(\alpha p - \frac{q}{p} \varepsilon)^{-1}$.

□

We are now in a position to prove Theorem 4.19.

Proof of Theorem 4.19. First note that

$$\nu_p |f - \nu_p f|^q \leq 2^q \nu_p |f - m|^q, \quad (4.21)$$

for all $m \in \mathbb{R}$. Now, for $R > 0$ and $L > 1$,

$$\begin{aligned} \nu_p |f - m|^q &= \nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \geq R\}}) + \nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \leq L\}}) \\ &\quad + \nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \geq L\}}). \end{aligned} \quad (4.22)$$

We treat each of the three terms of (4.22) separately.

First term of (4.22): This can be estimated using Lemma 4.20. Indeed

$$\begin{aligned} \nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \geq R\}}) &\leq \frac{1}{R} \nu_p (|f - m|^q N^{p-2} \|\cdot\|^2) \\ &\leq \frac{A}{R} \nu_p |\nabla_{\mathbb{G}} f|^q + \frac{B}{R} \nu_p |f - m|^q. \end{aligned} \quad (4.23)$$

Second term of (4.22): We have

$$\begin{aligned} \nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \leq L\}}) &\leq \nu_p (|f - m|^q \mathbf{1}_{\{N \leq L\}}) \\ &= \frac{1}{Z} \int_{\{N \leq L\}} |f(x) - m|^q e^{-\alpha N^p(x)} dx \\ &\leq \frac{1}{Z} \int_{\{N \leq L\}} |f(x) - m|^q dx. \end{aligned}$$

Take

$$m = \frac{1}{|B_{L_1}|} \int_{B_{L_1}} f(x) dx.$$

Then, since all homogeneous norms on \mathbb{G} are equivalent, by the Poincaré inequality in balls (see for example Theorem 5.6.1 of [23]) there exist constants P_0, L_1 such that

$$\nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \leq L\}}) \leq P_0 e^{\alpha L_1^p} \nu_p |\nabla_{\mathbb{G}} f|^q. \quad (4.24)$$

Third term of (4.22): Set $\bar{f} = f - m$ and $A_{L,R} := \{x \in \mathbb{G} : \|x\|^2 \leq R, N(x) \geq L\}$. Note that since $L > 1$ we have

$$\{x \in \mathbb{G} : \|x\|^2 N^{p-2}(x) \leq R, N(x) \geq L\} \subset A_{L,R}.$$

Thus

$$\nu_p(|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \geq L\}}) \leq \int_{A_{L,R}} |\bar{f}(x)|^q d\nu_p(x).$$

Recall that we can write $x = (w, z) \in \mathbb{G}$ for $w \in \mathbb{R}^n$ and $z \in \mathbb{R}^m$. For $e \in \{0, 1\}^m$, set

$$\mathcal{S}_e := \left\{ x = (w, z) \in \mathbb{G} : (-1)^{e_1} z_1 \geq 0, \dots, (-1)^{e_m} z_m \geq 0 \right\},$$

so that $\mathbb{G} = \cup_{e \in \{0,1\}^m} \mathcal{S}_e$. The reason for introducing these sets is so that in a particular \mathcal{S}_e , the signs of $z_j \in \mathbb{R}$ for $j \in \{1, \dots, m\}$ are known. By above, we then have

$$\nu_p(|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \geq L\}}) \leq \sum_{e \in \{0,1\}^m} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x)|^q d\nu_p(x). \quad (4.25)$$

We consider $\int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x)|^q d\nu_p(x)$ with $e = (0, \dots, 0)$ (the other cases are similar). Let $h \in \mathbb{G}$ be such that $\|h\| = 2\sqrt{R}$. Then we may write

$$\begin{aligned} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x)|^q d\nu_p(x) &\leq 2^{q-1} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x) - \bar{f}(xh)|^q d\nu_p(x) \\ &\quad + 2^{q-1} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q d\nu_p(x). \end{aligned} \quad (4.26)$$

Let $\gamma : [0, t] \rightarrow \mathbb{G}$ be a horizontal geodesic from 0 to h such that $|\dot{\gamma}(s)| \leq 1$ for $s \in [0, t]$. Then, by Hölder's inequality,

$$\begin{aligned} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x) - \bar{f}(xh)|^q d\nu_p(x) &= \int_{\mathcal{S}_e \cap A_{L,R}} \left| \int_0^t \frac{d}{ds} \bar{f}(x\gamma(s)) ds \right|^q d\nu_p(x) \\ &\leq t^{\frac{q}{p}} \int_0^t \int_{\mathcal{S}_e \cap A_{L,R}} |\nabla_{\mathbb{G}} f(x\gamma(s))|^q d\nu_p(x) ds. \end{aligned} \quad (4.27)$$

Using this estimate in (4.26), we arrive at

$$\begin{aligned} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x)|^q d\nu_p(x) &\leq 2^{q-1} d^{\frac{q}{p}}(h) \int_0^t \int_{\mathcal{S}_e \cap A_{L,R}} |\nabla_{\mathbb{G}} f(x\gamma(s))|^q d\nu_p(x) ds \\ &\quad + 2^{q-1} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q d\nu_p(x). \end{aligned} \quad (4.28)$$

Since we have chosen h such that $\|h\| = 2\sqrt{R}$, we have for $x \in A_{L,R}$

$$\|xh\| \geq \|h\| - \|x\| \geq 2\sqrt{R} - \sqrt{R} = \sqrt{R}. \quad (4.29)$$

We now claim that, for fixed R , we can choose h depending only on R , with $\|h\| = 2\sqrt{R}$, such that for large enough L

$$N^p(xh) \leq N^p(x), \quad \forall x \in \mathcal{S}_e \cap A_{L,R} \quad (4.30)$$

i.e. translation by h shifts points of $\mathcal{S}_e \cap A_{L,R}$ closer to the origin (with respect to the distance N).

Proof of claim (4.30): For $x = (w, z) \in \mathcal{S}_e \cap A_{L,R}$, we have

$$\|x\| = |w| \leq \sqrt{R}, \quad N(x) \geq L, \quad \text{and} \quad z_1 \geq 0, \dots, z_m \geq 0.$$

Let $h = (2\sqrt{R}, 0, \dots, 0, h_1, \dots, h_m) \in \mathbb{G} = \mathbb{R}^{n+m}$, for h_1, \dots, h_m only depending on R to be chosen later. Then, by the definition of the group law (see Theorem 2.2),

$$xh = \left(w_1 + 2\sqrt{R}, \dots, w_n, \right. \\ \left. z_1 + h_1 + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(1)} w_j \right), \dots, z_m + h_m + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(m)} w_j \right) \right),$$

so that

$$\begin{aligned} N^4(xh) - N^4(x) &= \left((w_1 + 2\sqrt{R})^2 + w_2^2 + \dots + w_n^2 \right)^2 + 16 \left(z_1 + h_1 + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(1)} w_j \right) \right)^2 \\ &\quad + \dots + 16 \left(z_m + h_m + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(m)} w_j \right) \right)^2 \\ &\quad - (w_1^2 + \dots + w_n^2)^2 - 16(z_1^2 + \dots + z_m^2). \end{aligned}$$

After expansion and cancellation, since we are taking x such that $\|x\| = |w| \leq \sqrt{R}$, we can bound all the remaining terms in the above expression that only involve w_1, \dots, w_n from above in terms of R . This will leave us with

$$\begin{aligned} N^4(xh) - N^4(x) &\leq K(R) + 32z_1 \left(h_1 + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(1)} w_j \right) \right) \\ &\quad + \dots + 32z_m \left(h_m + \sqrt{R} \left(\sum_{j=1}^n U_{1j}^{(m)} w_j \right) \right) \end{aligned}$$

for some constant K depending on R and the matrices $U^{(i)}$ for $i \in \{1, \dots, m\}$. Now, for $i \in \{1, \dots, m\}$ let $K_i(R)$ be the constant such that

$$\sqrt{R} \left| \sum_{j=1}^n U_{1j}^{(i)} w_j \right| \leq K_i(R)$$

for all $w \in \mathbb{R}^n$ such that $|w| \leq \sqrt{R}$ (so that K_i also depends on the matrix $U^{(i)}$). Then, since $z_i \geq 0$ for $i \in \{1, \dots, m\}$ by assumption, we have

$$N^4(xh) - N^4(x) \leq K(R) + 32z_1(h_1 + K_1(R)) + \dots + 32z_m(h_m + K_m(R)).$$

Let $\varepsilon > 0$, and take $h_i = -K_i(R) - \varepsilon$ for $i \in \{1, \dots, m\}$. Then

$$N^4(xh) - N^4(x) \leq K(R) - 32\varepsilon z_1 - \dots - 32\varepsilon z_m. \quad (4.31)$$

Now, since we are assuming that $N(x) \geq L$ and $|w| \leq \sqrt{R}$, it follows that

$$|z|^2 \geq \frac{1}{16}(L^4 - R^2).$$

Thus $z_j \geq \frac{1}{4\sqrt{m}}(L^4 - R^2)^{\frac{1}{2}}$ for at least one $j \in \{1, \dots, m\}$, so that by (4.31) we have

$$N^4(xh) - N^4(x) \leq K(R) - 8\varepsilon \frac{1}{\sqrt{m}}(L^4 - R^2)^{\frac{1}{2}}. \quad (4.32)$$

For big enough L the right-hand side of (4.32) is negative, which proves the claim (4.30).

We now use (4.29) and (4.30) to estimate the terms of (4.28). Indeed, using (4.29) we have that

$$\begin{aligned} 2^{q-1} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q d\nu_p(x) &\leq \frac{2^{q-1}}{R} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q \|xh\|^2 d\nu_p(x) \\ &\leq \frac{2^{q-1}}{R(L - N(h))^{p-2}} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q \|xh\|^2 N^{p-2}(xh) d\nu_p(x) \end{aligned} \quad (4.33)$$

for L large in comparison with $N(h)$. By (4.30) we also have

$$d\nu_p(x) = Z^{-1} e^{-\alpha N^p(x)} dx \leq Z^{-1} e^{-\alpha N^p(xh)} dx = d\nu_p(xh)$$

on $\mathcal{S}_e \cap A_{L,R}$, so that we can continue (4.33) to see that

$$2^{q-1} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q d\nu_p(x) \quad (4.34)$$

$$\begin{aligned} &\leq \frac{2^{q-1}}{R(L - N(h))^{p-2}} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(xh)|^q \|xh\|^2 N^{p-2}(xh) d\nu_p(xh) \\ &\leq \frac{2^{q-1}A}{R(L - N(h))^{p-2}} \nu_p |\nabla_{\mathbb{G}} f|^q + \frac{2^{q-1}B}{R(L - N(h))^{p-2}} \nu_p |f - m|^q \end{aligned} \quad (4.35)$$

where we have used the translational invariance of the Lebesgue measure, and Lemma 4.20 again.

For the first term of (4.28), note that there exists a constant $\tilde{K} = \tilde{K}(h)$ depending only on h (and hence only on R) such that

$$N^p(x\gamma(s)) - N^p(x) \leq \tilde{K}(h), \quad \forall x \in \mathcal{S}_e \cap A_{L,R}, \quad s \in [0, t].$$

This is because $N^p(x\gamma(s)) - N^p(x) \rightarrow 0$ as $N(x) \rightarrow \infty$ by the mean value theorem. Then

$$\begin{aligned} \int_0^t \int_{\mathcal{S}_e \cap A_{L,R}} |\nabla_{\mathbb{G}} f(x\gamma(s))|^q d\nu_p(x) ds &\leq e^{\tilde{K}(h)} \int_0^t \int_{\mathcal{S}_e \cap A_{L,R}} |\nabla_{\mathbb{G}} f(x\gamma(s))|^q d\nu_p(x\gamma(s)) ds \\ &\leq d(h) e^{\tilde{K}(h)} \nu_p |\nabla_{\mathbb{G}} f|^q. \end{aligned} \quad (4.36)$$

Using (4.36) together with (4.34) in (4.28) yields

$$\begin{aligned} \int_{\mathcal{S}_e \cap A_{L,R}} |\bar{f}(x)|^q d\nu_p(x) &\leq 2^{q-1} \left(d^{\frac{q}{p}+1}(h) e^{\tilde{K}(h)} + \frac{A}{R(L - N(h))^{p-2}} \right) \nu_p |\nabla_{\mathbb{G}} f|^q \\ &\quad + \frac{2^{q-1} B}{R(L - N(h))^{p-2}} \nu_p |f - m|^q. \end{aligned} \quad (4.37)$$

The key point is that the coefficient $\frac{2^{q-1} B}{R(L - N(h))^{p-2}}$ can be made as small as we wish by taking R large enough, provided L remains large in comparison. Although we have done the calculations for a specific $e \in \{0, 1\}^m$, the same may be done for arbitrary e (with a different choice of h). Thus, by (4.25), we see that there exist constants $C(R, L)$ and $\delta(R, L)$ such that

$$\nu_p (|f - m|^q \mathbf{1}_{\{\|\cdot\|^2 N^{p-2} \leq R\}} \mathbf{1}_{\{N \geq L\}}) \leq C(R, L) \nu_p |\nabla_{\mathbb{G}} f|^q + \delta(R, L) \nu_p |f - m|^q, \quad (4.38)$$

where $\delta(R, L)$ may be made as small as we wish by taking L and R large enough. This completes the estimate of the third term of (4.22).

It remains to insert the estimates (4.23), (4.24) and (4.38) into (4.22). Doing this we arrive at

$$\nu_p |f - m|^q \leq \left(\frac{A}{R} + P_0(L_1) e^{\alpha L^p} + C(R, L) \right) \nu_p |\nabla_{\mathbb{G}} f|^q + \left(\frac{B}{R} + \delta(R, L) \right) \nu_p |f - m|^q,$$

where R and L may be taken large enough so that $\frac{B}{R} + \delta(R, L) < 1$. Upon rearrangement, this inequality, combined with the observation (4.21), proves Theorem 4.19. \square

Conjecture 4.21. *Although our current techniques do not allow us to conclude that \mathcal{T}_p given by (4.10) has empty essential spectrum, we conjecture that this will be true for $p > 2$. This is a clear direction for further investigation.*

Acknowledgements The work of this paper came about as a result of some discussions with Prof. L. Saloff-Coste, Prof. A. Laptev and Prof. W. Hebisch to whom the author is very grateful.

References

- [1] D. BAKRY, F. BAUDOUIN, M. BONNEFONT, AND D. CHAFAÏ, *On gradient bounds for the heat kernel on the Heisenberg group*, J. Funct. Anal., 255 (2008), pp. 1905–1938.
- [2] D. BAKRY AND M. ÉMERY, *Hypercontractivité de semi-groupes de diffusion*, C. R. Acad. Sci. Paris Sér. I Math., 299 (1984), pp. 775 – 778.
- [3] F. BAUDOUIN, M. HAIRER, AND J. TEICHMANN, *Ornstein-Uhlenbeck processes on Lie groups*, J. Funct. Anal., 255 (2008), pp. 877–890.
- [4] A. BELLAÏCHE, *The tangent space in sub-Riemannian geometry*, in Sub-Riemannian Geometry, A. Bellaïche and J. J. Risler, eds., no. 144 in Progr. Math., Birkhäuser, 1996, pp. 1–78.
- [5] S. BOBKOV AND B. ZEGARLIŃSKI, *Entropy bounds and isoperimetry*, Mem. Amer. Math. Soc., 176 (2005).
- [6] A. BONFIGLIOLI, E. LANCONELLI, AND F. UGUZZONI, *Stratified Lie Groups and Potential Theory for their Sub-Laplacians*, Springer Monographs in Mathematics, Springer, 2007.
- [7] F. CIPRIANI, *Sobolev-Orlicz imbeddings, weak compactness, and spectrum*, J. Funct. Anal., 177 (2000), pp. 89–106.
- [8] B. DRIVER AND T. MELCHER, *Hypoelliptic heat kernel inequalities on the Heisenberg group*, J. Funct. Anal., 221 (2005), pp. 340–365.
- [9] N. ELDREDGE, *Precise estimates for the subelliptic heat kernel on H-type groups*, J. Math. Pures Appl., 92 (2009), pp. 52–85.
- [10] —, *Gradient estimates for the subelliptic heat kernel on H-type groups*, J. Funct. Anal., 258 (2010), pp. 504–533.
- [11] G. B. FOLLAND, *Subelliptic estimates and function spaces on nilpotent Lie groups*, Ark. Mat., 13 (1975), pp. 161–207.
- [12] G. GRILLO, *On Persson’s theorem in local Dirichlet spaces*, Z. Anal. Anwendungen, 17 (1998), pp. 329–338.

- [13] A. M. HANSSON AND A. LAPTEV, *Sharp spectral inequalities for the Heisenberg Laplacian*, in Groups and Analysis, K. Trent, ed., no. 354 in London Mathematical Society Lecture Note Series, CUP, 2008.
- [14] W. HEBISCH AND B. ZEGARLIŃSKI, *Coercive inequalities on metric measure spaces*, J. Funct. Anal., 258 (2010), pp. 814–851.
- [15] A. KAPLAN, *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc., 258 (1980), pp. 147–153.
- [16] L. D. LANDAU AND E. M. LIFSHITZ, *Quantum Mechanics: Non-relativistic Theory*, Pergamon Press, 1958.
- [17] H. Q. LI, *Estimation optimale du gradient du semi-groupe de la chaleur sur le groupe de Heisenberg*, J. Funct. Anal., 236 (2006), pp. 369–394.
- [18] T. MELCHER, *Hypoelliptic heat kernel inequalities on Lie groups*, PhD thesis, University of California, San Diego, 2004.
- [19] R. MONTI, *Some properties of Carnot-Carathéodory balls in the Heisenberg group*, Rend. Mat. Acc. Lincei, 11 (2000), pp. 155–167.
- [20] A. PERSSON, *Bounds for the discrete part of the spectrum of a semibounded Schrödinger operator*, Math. Scand., 8 (1960), pp. 143–153.
- [21] M. REED AND B. SIMON, *IV: Analysis of Operators*, Methods of Modern Mathematical Physics, Academic Press, 1978.
- [22] L. C. G. ROGERS AND D. WILLIAMS, *Diffusions, Markov Processes, and Martingales: Volume 1, Foundations*, Cambridge Mathematical Library, Cambridge University Press, 2000.
- [23] L. SALOFF-COSTE, *Aspects of Sobolev-Type Inequalities*, no. 289 in London Mathematical Society Lecture Note Series, Cambridge University Press, 2002.
- [24] N. T. VAROPOULOS, L. SALOFF-COSTE, AND T. COULHON, *Analysis and Geometry on Groups*, no. 100 in Cambridge Tracts in Mathematics, Cambridge University Press, 1992.
- [25] F. Y. WANG, *Functional inequalities for empty essential spectrum*, J. Funct. Anal., 170 (2000), pp. 219–245.
- [26] —, *Functional inequalities and spectrum estimates: the infinite measure case*, J. Funct. Anal., 194 (2002), pp. 288–310.

- [27] —, *Super and weak Poincaré inequalities for hypoelliptic operators*, Acta Math. Appl. Sin. Engl. Ser., 25 (2009), pp. 617–630.
- [28] P.-A. ZITT, *Super Poincaré inequalities, Orlicz norms and essential spectrum*. Preprint, available at hal-00426367, 2010.