

Stability analysis of a Vlasov-Wave system describing particles interacting with their environment

Stephan De Bièvre¹

*Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlevé, F-59000 Lille, France
‡ Equipe-Projet MEPHYSTO, Centre de Recherche INRIA Lille Nord Europe,
Parc Scientifique de la Haute Borne, 40, avenue Halley B.P. 70478
F-59658 Villeneuve d'Ascq cedex, France*

Thierry Goudon

*Université Côte d'Azur, Inria, CNRS, LJAD,
Parc Valrose, F-06108 Nice, France*

Arthur Vasseur

*Université Côte d'Azur, Inria, CNRS, LJAD,
‡ Basque Center for Applied Mathematics
Alameda de Mazarredo, 14
E-48009 Bilbao, Basque Country - Spain*

Abstract

We study a kinetic equation of the Vlasov-Wave type, which arises in the description of the behaviour of a large number of particles interacting weakly with an environment, composed of an infinite collection of local vibrational degrees of freedom, modeled by wave equations. We use variational techniques to establish the existence of large families of stationary states for this system, and analyze their stability.

Keywords:

Vlasov-like equations, Interacting particles, Dynamical stability, Inelastic Lorentz gas

2000 MSC: 82C70, 70F45, 37K05, 74A25

Email addresses: stephan.de-bievre@univ-lille1.fr (Stephan De Bièvre),
thierry.goudon@inria.fr (Thierry Goudon), arthur.vasseur@unice.fr (Arthur Vasseur)

¹S.D.B. acknowledges the support of the Labex CEMPI (ANR-11-LABX-0007-01)

1. Introduction

This paper is concerned with the following PDE system

$$\partial_t f + v \cdot \nabla_x f - \nabla_x(V + \Phi) \cdot \nabla_v f = 0, \quad t \geq 0, x \in \mathbb{R}^d, v \in \mathbb{R}^d, \quad (1a)$$

$$(\partial_{tt}^2 \Psi - c^2 \Delta_z \Psi)(t, x, z) = -\sigma_2(z) \int_{\mathbb{R}^d} \sigma_1(x - y) \rho(t, y) dy, \quad (1b)$$

$$t \geq 0, x \in \mathbb{R}^d, z \in \mathbb{R}^n,$$

$$\rho(t, x) = \int_{\mathbb{R}^d} f(t, x, v) dv, \quad (1c)$$

$$\Phi(t, x) = \iint_{\mathbb{R}^d \times \mathbb{R}^n} \Psi(t, y, z) \sigma_1(x - y) \sigma_2(z) dz dy, \quad t \geq 0, x \in \mathbb{R}^d. \quad (1d)$$

This system was introduced in [8, 26], where also the existence of solutions was established and some asymptotic issues treated. In particular, a certain asymptotic regime brings out a quite unexpected connection between this system and the gravitational Vlasov–Poisson system. In this work, we wish to address a first aspect of the long-time behaviour of the system by investigating the existence of stationary solutions and their stability.

We recall the system (1a)–(1d) describes the interaction between a large set of particles and their environment. The particles are described by their distribution function in phase space: $f(t, x, v)$, where $t \geq 0$, $x \in \mathbb{R}^d$, $v \in \mathbb{R}^d$ are the time, space and velocity variables, respectively. More precisely, we interpret this quantity so that, given $\mathcal{O} \times \mathcal{V} \subset \mathbb{R}^d \times \mathbb{R}^d$, $\iint_{\mathcal{O} \times \mathcal{V}} f(t, x, v) dv dx$ gives the mass of particles having at time $t \geq 0$ their position in \mathcal{O} and their velocity in \mathcal{V} . The environment is seen as a vibrating medium, the state of which is described by the function Ψ . Note that the vibrations take place in the direction $z \in \mathbb{R}^n$, which is *transverse* to the particles' motion; they are characterized by the wave speed $c > 0$. The coupling between the particles and the environment is embodied in the form functions σ_1, σ_2 , which are both non negative, smooth and radially symmetric functions. Finally, the particles are also subjected to a confining potential V . We shall give details later on the technical assumptions and start by discussing the modelling issues.

The system (1a)–(1d) is a kinetic version of a model introduced by L. Bruneau and S. De Bièvre [5] for a *single* classical particle interacting with its environment. It is useful to keep in mind the picture proposed in

[5] of a particle moving through an infinite set of n -dimensional membranes, one for each $x \in \mathbb{R}^d$. Denoting by $t \mapsto q(t)$ the position of the particle, the dynamic of such a particle is described by the following set of differential equations

$$\begin{cases} \ddot{q}(t) = -\nabla V(q(t)) - \iint_{\mathbb{R}^d \times \mathbb{R}^n} \sigma_1(q(t) - y) \sigma_2(z) \nabla_x \Psi(t, y, z) dy dz, \\ \partial_{tt}^2 \Psi(t, x, z) - c^2 \Delta_z \Psi(t, x, z) = -\sigma_2(z) \sigma_1(x - q(t)), \quad x \in \mathbb{R}^d, z \in \mathbb{R}^n. \end{cases} \quad (2)$$

For the system (2), due to the energy exchanges between the particles and the environment, and the evacuation of energy in the membranes, the coupling eventually behaves like a friction force on the particle so that the particle asymptotically stops at the minimum of the confining potential V . Further situations are dealt with in [5], and we also refer the reader to [1, 9, 10, 11, 21, 25] for thorough numerical experiments and analytical studies on (2) and related models. The system (2) can be generalized by considering a set of N particles, and the mean field regime $N \rightarrow \infty$ leads to (1a)–(1d), see [17, 26].

In [2], hypocoercivity techniques are adapted from [13] to investigate the large time behavior of the solutions, when the kinetic equation (1a) additionally contains the Fokker–Planck operator $\nabla_v \cdot (vf + \nabla_v f)$. This operator induces some dissipative effects in the model which simplifies the asymptotic analysis. However, the stationary solution

$$\mathcal{M}_{\text{eq}}(x, v) = Z_{\text{eq}} \exp\left(-\frac{|v|^2}{2} - V(x) - \Phi_{\text{eq}}(x)\right) \quad (3)$$

exhibited in [2, 26] defines also a stationary solution of (1a)–(1d). It is then natural to investigate the stability of this solution, a task we turn to in this paper. In (3), Φ_{eq} is implicitly defined by the non-linear equation

$$\Phi_{\text{eq}}(x) = -\frac{\kappa}{c^2} \sigma_1 \star \sigma_1 \star \int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(x, v) dv, \quad \kappa = \int_{\mathbb{R}^n} \frac{|\widehat{\sigma}_2(\xi)|^2}{|\xi|^2} d\xi > 0. \quad (4)$$

Note that the definition of κ requires $n \geq 3$. It can be equivalently expressed as

$$\kappa = \int_{\mathbb{R}^n} |\nabla \Gamma(z)|^2 dz > 0,$$

with Γ the solution of

$$\Delta \Gamma(z) = \sigma_2(z), \quad z \in \mathbb{R}^n.$$

Equations (3) and (4) define a family of stationary solutions of (1a)–(1d), parametrized here by $Z_{\text{eq}} \geq 0$. Since (1a) conserves the total mass, we address the question of the stability of the stationary solution which has the same mass as the initial data, namely Z_{eq} is the normalizing constant such that

$$\begin{aligned} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \mathcal{M}_{\text{eq}}(x, v) \, dv \, dx &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(0, x, v) \, dv \, dx \\ &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(t, x, v) \, dv \, dx. \end{aligned}$$

In [2, 26] this solution is shown to be *linearly stable* for the collisionless system (1a)–(1d). We wish to investigate further the issue of the existence of these and other stationary solutions and their non-linear stability. We point out that the large time behavior for the kinetic model (1a)–(1d), or even for the N -particle version of (2), differs considerably from the situation investigated in [5] for a single particle, precisely because of the intricate interactions between all the particles when they are close to each other. When dealing with a single particle, with V a confining potential as considered here, the asymptotic behavior is as expected from basic physical intuition. Assuming for simplicity the potential has a unique critical point, which is its absolute minimum, it was shown the particle comes (exponentially fast) to rest at this minimum, see [5, Theorem 4]. This is – intuitively at least – a consequence of the fact that, as long as the particle moves, it will loose energy to the membranes, so that it has to come to a full stop. Which it can only do where no external force is applied by the external potential. With several particles, the underlying physics is more complex since when two particles are close enough to each other, they activate the membranes in a common neighborhood, which could lead to more complicated energy exchanges between the particle and the membranes. So, while a stationary solution with all particles at the minimum of the potential still exists, it is thus not clear if other stationary solutions may or may not exist. For the kinetic model, which corresponds to the statistical description of many particles in a mean field regime when the coupling between the particles is weak, we shall see that the model admits infinitely many stationary solutions with different shape.

In this paper we thus go back to the stationary solutions of the Vlasov-Wave system (1a)–(1d). Using the Hamiltonian structure at the basis of the discussion in [5], and the conservation of the Casimir–functionals of the Vlasov equation (1a), we develop a variational approach, strongly inspired by the analysis of the gravitational Vlasov–Poisson system [14, 18, 19, 24, 27];

see also [3, 6] for the electrostatic Vlasov–Poisson system.

In Section 2, we construct an infinite family of stationary solutions $(f_{\text{eq}}, \Phi_{\text{eq}})$ of the system (1a)–(1d) through the minimization of suitable energy functionals, which are constants of the motion of the system, under a mass constraint. Section 3 investigates the dynamical stability of these stationary solutions. We conclude this Introduction with a rough formulation of our main result, giving a flavour of it. (See Theorem 3.2 for a precise statement.) Each energy–Casimir functional defines an equilibrium $(f_{\text{eq}}, \Phi_{\text{eq}})$ and a natural measure, that we denote by $(f, \Phi) \mapsto \mathcal{D}(f, \Phi | f_{\text{eq}}, \Phi_{\text{eq}})$, which evaluates how far a pair (f, Φ) is from this equilibrium. The stability statement then says that, if the initial data is close enough to the equilibrium state, then it remains close to it forever: for any $\epsilon > 0$, there exists $\eta > 0$ such that $\mathcal{D}(f(0), \Phi(0) | f_{\text{eq}}, \Phi_{\text{eq}}) \leq \eta$ implies $\mathcal{D}(f(t), \Phi(t) | f_{\text{eq}}, \Phi_{\text{eq}}) \leq \epsilon$ for any $t \geq 0$. As we shall see below, beyond a few technical assumptions, the proof requires $n \geq 3$ and the wave speed c to be large enough, which both seem to be important assumptions on physical grounds.

Whether analogous stationary states also exist in the original N -particle model is a difficult open question that we do not address here and that is related to the question of the time-scale on which the Vlasov-Wave equation correctly describes the N -particle model. The asymptotic trend to the equilibrium, in absence of collisional mechanisms, is also a delicate open issue.

2. Stationary solutions

We start our study with a few formal manipulations, which will motivate our choice of functional framework for the rigorous study of the stationary solutions of (1a)–(1d) that is provided in the second part of this section. A first crucial observation is that (1a)–(1d) conserves an energy functional that encodes the energy exchanges between the particles and their environment:

$$\begin{aligned} \mathcal{E}(f, \Psi, \pi) = & \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(\frac{|v|^2}{2} + V(x) \right) f \, dv \, dx + \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi[\Psi] f \, dv \, dx \\ & + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} (|\pi|^2 + c^2 |\nabla_z \Psi|^2) \, dy \, dx. \end{aligned} \tag{5}$$

Here we have used the following notation: for a function $\Psi : \mathbb{R}^d \times \mathbb{R}^n \rightarrow \mathbb{R}$,

$$\Phi[\Psi](x) = \int_{\mathbb{R}^d} \sigma_1(x-y) \left(\int_{\mathbb{R}^n} \sigma_2(z) \Psi(y, z) \, dz \right) \, dy. \tag{6}$$

Then, we have energy conservation in the sense that the solutions of (1a)–(1d) satisfy

$$\mathcal{E}(f, \Psi, \partial_t \Psi)(t) = \mathcal{E}(f, \Psi, \partial_t \Psi)(0). \quad (7)$$

Of course, the particles’ total mass is also conserved

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f(t, x, v) \, dv \, dx = \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(0, x, v) \, dv \, dx. \quad (8)$$

Finally, since the field $(x, v) \mapsto (v, -\nabla_x V(x) - \nabla_x \Phi(t, x))$ is divergence-free, equation (1a) also conserves any non-linear functional (the “Casimir-functionals”)

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} h(f(t, x, v)) \, dv \, dx = \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(f(0, x, v)) \, dv \, dx, \quad (9)$$

for a given function $h : [0, \infty) \rightarrow \mathbb{R}$.

This suggests to search for stationary solutions of (1a)–(1d) as minimizers of a Lyapounov functional using (7), (8) and (9). The natural energy (5) does not have enough structure to serve as a Lyapounov functional for the evolution problem. The idea is thus to combine (5) and (9) and to study the critical points of this Casimir-energy functional, under a mass constraint. It turns out that the critical points are also stationary solutions of (1a)–(1d). Moreover, the minimization property defines the framework that allows us to investigate the stability issue. We refer for instance the reader to [3, 6, 12, 18, 27] for similar reasonings, see also the overview [24]. We proceed as follows. Let $h : [0, \infty) \rightarrow \mathbb{R}$ be a *strictly convex* function. We set

$$\mathcal{J}(f, \Psi, \pi) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(f) \, dv \, dx + \mathcal{E}(f, \Psi, \pi).$$

Let us denote by m the mass of the initial particle distribution function

$$m = \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(0, x, v) \, dv \, dx.$$

We wish to define $(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}})$ as the minimizer of \mathcal{J} under the constraint

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f(x, v) \, dv \, dx = m.$$

We thus introduce, for $\lambda \in \mathbb{R}$, $\mathcal{J}_\lambda = \mathcal{J} + \lambda(\iint_{\mathbb{R}^d \times \mathbb{R}^d} f \, dv \, dx - m)$. The Euler–Lagrange relations yield the following conclusions

- with $\partial_\pi \mathcal{J}_\lambda$ we simply get $\pi_{\text{eq}} = 0$,

- with $\partial_\Psi \mathcal{J}_\lambda$ we obtain

$$\sigma_2(z) \left(\sigma_1 \star \int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(\cdot, v) \, dv \right) (x) = c^2 \Delta_z \Psi_{\text{eq}}(x, z).$$

Denoting by Γ the solution of $\Delta_z \Gamma = \sigma_2$, we are led to

$$\Psi_{\text{eq}}(x, z) = \frac{\Gamma(z)}{c^2} \sigma_1 \star \left(\int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(\cdot, v) \, dv \right) (x).$$

The associated equilibrium potential thus reads

$$\begin{aligned} \Phi_{\text{eq}}(x) &= \int_{\mathbb{R}^d} \sigma_1(x-y) \left(\int_{\mathbb{R}^n} \sigma_2(z) \Psi_{\text{eq}}(y, z) \, dz \right) \, dy \\ &= \sigma_1 \star \sigma_1 \star \left(\int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(\cdot, v) \, dv \right) (x) \times \left(-\frac{1}{c^2} \int_{\mathbb{R}^n} |\nabla_z \Gamma(z)|^2 \, dz \right) \\ &= -\frac{\kappa}{c^2} \sigma_1 \star \sigma_1 \star \left(\int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(\cdot, v) \, dv \right) (x). \end{aligned}$$

- with $\partial_f \mathcal{J}_\lambda$ we infer

$$\begin{aligned} h'(\mathcal{M}_{\text{eq}}(x, v)) &= -\frac{|v|^2}{2} - V(x) - \sigma_1 \star \int_{\mathbb{R}^n} \sigma_2(z) \Psi_{\text{eq}}(\cdot, z) \, dz - \lambda_{\text{eq}} \\ &= -\frac{|v|^2}{2} - V(x) \\ &\quad + \frac{\kappa}{c^2} \sigma_1 \star \sigma_1 \star \left(\int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(\cdot, v) \, dv \right) (x) - \lambda_{\text{eq}}, \end{aligned}$$

with λ_{eq} the Lagrange multiplier associated to the mass constraint.

Therefore, the equilibrium distribution function is defined by the implicit formula

$$\mathcal{M}_{\text{eq}}(x, v) = \Upsilon \left(\frac{|v|^2}{2} + V(x) - \frac{\kappa}{c^2} \Sigma \star \rho_{\text{eq}}(x) + \lambda_{\text{eq}} \right),$$

$$\Sigma = \sigma_1 \star \sigma_1, \quad \rho_{\text{eq}} = \int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}} \, dv,$$

where

$$\Upsilon(s) = (h')^{-1}(-s),$$

and the Lagrange multiplier λ_{eq} is determined by

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \mathcal{M}_{\text{eq}} \, dv \, dx = m.$$

We readily check that

$$(v \cdot \nabla_x - \nabla_x(V + \Psi_{\text{eq}}) \cdot \nabla_v) \mathcal{M}_{\text{eq}} = 0$$

so that \mathcal{M}_{eq} is a stationary solution of (1a), as anticipated. This comes from the fact that, writing $\mathcal{E}_{\text{eq}}(x, v) = \frac{|v|^2}{2} + V(x) - \frac{\kappa}{c^2} \Sigma \star \rho_{\text{eq}}(x) = \frac{|v|^2}{2} + V(x) + \Phi_{\text{eq}}(x)$, \mathcal{M}_{eq} can be interpreted as a function of this energy and $(v \cdot \nabla_x - \nabla_x(V + \Psi_{\text{eq}}) \cdot \nabla_v) \mathcal{M}_{\text{eq}}$ is the Lie bracket of \mathcal{E}_{eq} and \mathcal{M}_{eq} . Moreover, Ψ_{eq} is a stationary solution of the associated wave equation (1b)–(1c) while finally Φ_{eq} satisfies (1d).

The specific case $h(s) = s \ln(s) - s$ leads to (3), with $\Upsilon(s) = e^{-s}$, $Z_{\text{eq}} = e^{-\lambda_{\text{eq}}}$. With the change of variable $v = \sqrt{2s} \, \omega$, $s \geq 0$, $\omega \in \mathbb{S}^{d-1}$, we can rewrite

$$\begin{aligned} \rho_{\text{eq}}(x) &= \mu(V(x) + \Phi_{\text{eq}}(x) + \lambda_{\text{eq}}), & \Phi_{\text{eq}} &= -\frac{\kappa}{c^2} \Sigma \star \rho_{\text{eq}}, \\ \mu(U) &= 2^{d/2-1} |\mathbb{S}^{d-1}| \int_0^\infty s^{d/2-1} \Upsilon(s + U) \, ds. \end{aligned}$$

Through this approach we have therefore defined a broad class of stationary states for (1a)–(1d), in contrast to the case of a single particle where, as mentioned before, a unique equilibrium exists. We now investigate their stability.

Let us now detail the technical requirements on which the analysis is based. Throughout the paper we assume

- a) $n \geq 3$,
- b) σ_1, σ_2 are C^∞ , compactly supported, radially symmetric and non negative.

Assumption a) already appears in [5] as well as in [8]. In particular, we should bear in mind that the definition of κ uses this assumption. We refer the reader to [5] for further comments on the physical motivation for assuming $n \geq 3$. Assumption b) is certainly far from optimal, as will become clear from the discussion below, but it stays within the framework introduced and motivated in [5]. To give a precise definition of the coupling term in the

energy (5), we proceed as follows. First, we define, for $\Psi : \mathbb{R}^d \times \mathbb{R}^n \rightarrow \mathbb{R}$, and $\rho : \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\mathcal{B}(\Psi, \rho) = \int_{\mathbb{R}^d} (\sigma_1 \star \rho)(x) \left(\int_{\mathbb{R}^n} \sigma_2(z) \Psi(x, z) dz \right) dx.$$

Then one has

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi[\Psi] f dv dx = \mathcal{B}(\Psi, \rho),$$

where $\Phi[\Psi]$ is defined in (6).

Lemma 2.1. *There exists a constant C , which does not depend on ψ, ρ , such that*

$$|\mathcal{B}(\Psi, \rho)| \leq C \|\sigma_2\|_{L^{2n/(n+2)}(\mathbb{R}^n)} \|\sigma_1\|_{L^2(\mathbb{R}^d)} \|\nabla_z \Psi\|_{L^2(\mathbb{R}^n \times \mathbb{R}^d)} \|\rho\|_{L^1(\mathbb{R}^d)}.$$

Proof. We shall use the Gagliardo-Nirenberg-Sobolev inequality, see e.g. [23, p. 125], [4, Theorem IX.9],

$$\|u\|_{L^{2n/(n-2)}(\mathbb{R}^n)} \leq C \|\nabla u\|_{L^2(\mathbb{R}^n)}.$$

Then, we get

$$\begin{aligned} |\mathcal{B}(\Psi, \rho)| &\leq \left(\int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^n} \sigma_2(z) \Psi(x, z) dz \right|^2 dx \right)^{1/2} \left(\int_{\mathbb{R}^d} |\sigma_1 \star \rho(x)|^2 dx \right)^{1/2} \\ &\leq \left(\int_{\mathbb{R}^d} \|\sigma_2\|_{L^{2n/(n+2)}(\mathbb{R}^n)}^2 \left(\int_{\mathbb{R}^n} |\Psi(x, z)|^{2n/(n-2)} dz \right)^{(n-2)/n} dx \right)^{1/2} \\ &\quad \times \|\sigma_1\|_{L^2(\mathbb{R}^d)} \|\rho\|_{L^1(\mathbb{R}^d)} \\ &\leq \|\sigma_2\|_{L^{2n/(n+2)}(\mathbb{R}^n)} \left(\int_{\mathbb{R}^d} C^2 \left(\int_{\mathbb{R}^n} |\nabla_z \Psi(x, z)|^2 dz \right) dx \right)^{1/2} \\ &\quad \times \|\sigma_1\|_{L^2(\mathbb{R}^d)} \|\rho\|_{L^1(\mathbb{R}^d)}. \end{aligned}$$

■

We assume that the following requirements are fulfilled:

- (H0) For any $x \in \mathbb{R}^d$, we have $V(x) \geq 0$, and $\lim_{|x| \rightarrow \infty} V(x) = +\infty$.
- (H1) Either $h(s) = s \ln(s) - s$ or $h : [0, \infty) \rightarrow [0, \infty)$ is a strictly convex C^2 function such that

$$h(0) = 0 \text{ and } \lim_{s \rightarrow \infty} \frac{h(s)}{s} = +\infty,$$

with h' defined from $[0, \infty)$ to $[\eta_0, +\infty)$ with $\eta_0 \geq 0$. (That η_0 is non negative follows from the assumptions: $h(s)$ is non negative and vanishes for $s = 0$.)

(H2) Let us define

$$\Upsilon : s \in \mathbb{R} \longmapsto \Upsilon(s) = \begin{cases} 0 & \text{if } s \geq -\eta_0, \\ (h')^{-1}(-s) & \text{if } -\infty < s \leq -\eta_0. \end{cases}$$

For any $a \in \mathbb{R}$, we have

$$\Upsilon\left(\frac{|v|^2}{2} + V(x) + a\right) \in L^1(\mathbb{R}^d \times \mathbb{R}^d).$$

For $h(s) = s \ln(s) - s$, we further assume $x \mapsto e^{-\alpha V(x)} \in L^1(\mathbb{R}^d)$ for some $0 < \alpha < 1/4$.

(H3) For any compact set $K \subset \mathbb{R}$, we can find a function $\mathcal{Z}_K \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$ such that for any $a \in K$, we have $|\Upsilon\left(\frac{|v|^2}{2} + V(x) + a\right)| \leq \mathcal{Z}_K(x, v)$.

Assumptions **(H0)** is a confining assumption on the external potential, strengthened by the integrability condition in **(H2)**. It is likely that these assumptions on h , that govern the shape of the equilibrium state, can be generalized, as will become apparent in the proof. With $h(s) = s \ln(s) - s$, we obtain the equilibrium function devised in [2], which is proportional to $\exp\left(-\frac{|v|^2}{2} - V(x) - \Phi_{\text{eq}}(x)\right)$, see (3), the self-consistent potential Φ_{eq} being determined by an integral equation. Another relevant example is given by $h(s) = s^p$, for $p > 1$. Then, the equilibrium function casts as

$$\mathcal{M}_{\text{eq}}(x, v) = \left[-\frac{1}{p} \left(\frac{|v|^2}{2} + V(x) + \Phi_{\text{eq}}(x) + \lambda_{\text{eq}} \right) \right]_+^{1/(p-1)},$$

where the equilibrium potential Φ_{eq} is still determined by an integral equation and $\lambda_{\text{eq}} \in \mathbb{R}$ ensures the mass constraint. We shall see that $x \mapsto \Phi_{\text{eq}}(x)$ is bounded; hence, due to **(H0)**, we note that this equilibrium function is compactly supported.

Lemma 2.2. *Suppose **(H0)**–**(H2)**. Let $m > 0$. For any function Φ continuous and bounded on \mathbb{R}^d , we can find $\lambda[\Phi] \in \mathbb{R}$ such that*

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) dv dx = m.$$

Furthermore, for $c > 0$, let us introduce the set

$$\mathcal{C} = \left\{ \Phi \in C^0(\mathbb{R}^d), -\frac{\kappa m}{c^2} \|\Sigma\|_{L^\infty(\mathbb{R}^d)} \leq \Phi(x) \leq 0 \text{ for all } x \in \mathbb{R}^d \right\}.$$

There exists $\mathcal{K}_1(c) > 0$, such that $|\lambda[\Phi]| \leq \mathcal{K}_1(c)$ holds for any $\Phi \in \mathcal{C}$.

Proof. The function $s \mapsto \Upsilon(s)$ is strictly decreasing from $(-\infty, -\eta_0)$ to $(0, +\infty)$ (but it is defined from \mathbb{R} to $[0, \infty)$). For Φ bounded and continuous, by **(H0)**–**(H2)**, the mapping

$$\mathbb{M} : \lambda \mapsto \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda\right) dv dx$$

is well defined, decreasing, continuous (by applying Lebesgue's theorem), and it satisfies

$$\lim_{\lambda \rightarrow +\infty} \mathbb{M}(\lambda) = 0, \quad \lim_{\lambda \rightarrow -\infty} \mathbb{M}(\lambda) = +\infty.$$

The conclusion follows directly by the intermediate value theorem.

The bound on $\Phi \mapsto \lambda[\Phi]$ follows from the following argument. According to **(H1)**–**(H2)**, $s \mapsto \Upsilon(s)$ is non increasing and we get, for any $\Phi \in \mathcal{C}$,

$$\begin{aligned} & \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) + \lambda[\Phi]\right) dv dx \\ & \leq \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) dv dx = m \\ & \leq \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) - \frac{\kappa}{c^2} \|\Sigma\|_{L^\infty(\mathbb{R}^d)} + \lambda[\Phi]\right) dv dx. \end{aligned}$$

If $\lambda[\Phi] \rightarrow +\infty$ (resp. $\lambda[\Phi] \rightarrow -\infty$), the bound from above tends to 0 (resp. the bound from below tends to $+\infty$), which leads to a contradiction. ■

Having disposed of these preliminaries, we turn to the analysis of the equation satisfied by the critical points of the functional \mathcal{J} on the mass shell.

Proposition 2.3. *Let $m > 0$. Suppose **(H0)**–**(H3)**. There exists $c_0 > 0$ such that for any $c > c_0$, the equations*

$$\begin{aligned} \mathcal{M}_{\text{eq}}(x, v) &= \Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi_{\text{eq}}(x) + \lambda_{\text{eq}}\right), \\ \Phi_{\text{eq}}(x) &= -\frac{\kappa}{c^2} \Sigma \star \rho_{\text{eq}}, \quad \rho_{\text{eq}}(x) = \int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}}(x, v) dv, \\ \iint_{\mathbb{R}^d \times \mathbb{R}^d} \mathcal{M}_{\text{eq}}(x, v) dv dx &= \int_{\mathbb{R}^d} \rho_{\text{eq}}(x) dx = m, \end{aligned}$$

for the critical points of \mathcal{J} with mass constraint admit a unique solution.

Proof. We consider the map

$$\mathcal{T} : \Phi \in \mathcal{C} \mapsto \mathcal{T}[\Phi] = -\frac{\kappa}{c^2} \Sigma \star \rho[\Phi],$$

where

$$\rho[\Phi](x) = \int_{\mathbb{R}^d} \Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) dv$$

and $\lambda[\Phi] \in \mathbb{R}$ is such that

$$\int_{\mathbb{R}^d} \rho[\Phi](x) dx = m.$$

Lemma 2.2 ensures that this definition makes sense. In fact, a direct application of the implicit function theorem (that uses **(H3)**) tells us that the mapping $\Phi \mapsto \lambda[\Phi]$ is C^1 and we are going to compute its derivative. Moreover, \mathcal{T} maps \mathcal{C} into itself. We see Φ_{eq} as a fixed point of \mathcal{T} and we shall prove that \mathcal{T} is a contraction provided c is large enough. Let Φ and Φ' in \mathcal{C} . We start with the simple observation

$$\|\mathcal{T}[\Phi] - \mathcal{T}[\Phi']\| \leq \frac{\kappa}{c^2} \|\Sigma\|_{L^\infty(\mathbb{R}^d)} \|\rho[\Phi] - \rho[\Phi']\|_{L^1(\mathbb{R}^d)}.$$

Writing $\Phi_\theta = \Phi + \theta(\Phi' - \Phi)$ and $\lambda_\theta = \lambda[\Phi_\theta]$ for $0 \leq \theta \leq 1$, we obtain

$$\begin{aligned} |\rho[\Phi] - \rho[\Phi']|(x) &= \left| \int_0^1 \int_{\mathbb{R}^d} \frac{d}{d\theta} \left[\Upsilon\left(\frac{|v|^2}{2} + V(x) + \Phi_\theta(x) + \lambda_\theta\right) \right] dv d\theta \right| \\ &= \left| \int_0^1 \int_{\mathbb{R}^d} \Upsilon'\left(\frac{|v|^2}{2} + V(x) + \Phi_\theta(x) + \lambda_\theta\right) \right. \\ &\quad \left. \times \left\{ (\Phi' - \Phi)(x) + \lambda'[\Phi_\theta](\Phi' - \Phi) \right\} d\theta dv \right|. \end{aligned}$$

We find the (signed) measure $\lambda'[\Phi]$ by taking the derivative with respect to Φ of the mass constraint: for any continuous and bounded function φ , we get

$$0 = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon'\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) (\varphi(x) + \lambda'[\Phi](\varphi)) dv dx,$$

the formula makes sense owing to **(H3)**. Therefore, we obtain

$$\lambda'[\Phi](\varphi) = \frac{- \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon'\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) \varphi(x) dv dx}{\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon'\left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi]\right) dv dx}.$$

We deduce from this relation that

$$|\lambda'[\Phi](\varphi)| \leq \|\varphi\|_{L^\infty(\mathbb{R}^d)}. \quad (10)$$

Let us temporarily assume that for some $c_0 > 0$, we can find a constant $\mathcal{K}(c_0)$ such that for any $c \geq c_0$ and $\Phi \in \mathcal{C}$ we have

$$\left| \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon' \left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi] \right) dv dx \right| \leq \mathcal{K}(c_0). \quad (11)$$

Owing to (11), it follows that

$$|\rho[\Phi] - \rho[\Phi']|(x) \leq 2\mathcal{K}(c_0)\|\Phi - \Phi'\|_{L^\infty(\mathbb{R}^d)}$$

holds for any $\Phi, \Phi' \in \mathcal{C}$ and $c \geq c_0$. We conclude that

$$|\mathcal{T}[\Phi] - \mathcal{T}[\Phi']| \leq \frac{2\kappa}{c^2} \mathcal{K}(c_0) \|\Sigma\|_{L^\infty(\mathbb{R}^d)} \|\Phi - \Phi'\|_{L^\infty(\mathbb{R}^d)}.$$

We now prove (11). As $c \rightarrow \infty$ the functions in \mathcal{C} converge uniformly to 0; moreover, by virtue of (10), we see that $\lambda[\Phi]$ tends to $\lambda_0 \in \mathbb{R}$, which is determined by

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon \left(\frac{|v|^2}{2} + V(x) + \lambda_0 \right) dv dx = m,$$

see Lemma 2.2. For $c > 0$ and $\Phi \in \mathcal{C}$, it follows from **(H3)** that the integral

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon' \left(\frac{|v|^2}{2} + V(x) + \Phi(x) + \lambda[\Phi] \right) dv dx$$

is well defined. In addition, as $c \rightarrow \infty$ it tends to

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \Upsilon' \left(\frac{|v|^2}{2} + V(x) + \lambda_0 \right) dv dx$$

by continuity and the dominated convergence theorem. It allows us to check that (11) holds for $c \geq c_0$ large enough. Moreover we have

$$\lim_{c \rightarrow \infty} \frac{\mathcal{K}(c_0)}{c^2} = 0,$$

and c_0 can be chosen large enough so that \mathcal{T} is a contraction on \mathcal{C} for $c \geq c_0$. \blacksquare

We turn to the minimization problem. We extend h by $+\infty$ when its argument is negative, so that \mathcal{J} is defined over

$$\begin{aligned} f &\in L^1(\mathbb{R}^d \times \mathbb{R}^d), & (|v|^2 + V(x))f &\in L^1(\mathbb{R}^d \times \mathbb{R}^d), \\ \nabla_z \Psi &\in L^2(\mathbb{R}^d \times \mathbb{R}^n), & \pi &\in L^2(\mathbb{R}^d \times \mathbb{R}^n). \end{aligned} \quad (12)$$

We wish to minimize \mathcal{J} under a mass constraint; the connection with the

critical points discussed in Proposition 2.3 is established through the following claim.

Lemma 2.4. *Let $m > 0$. If \mathcal{J} admits a minimizer (f, Ψ, π) over the functions in (12) satisfying the mass constraint $\iint_{\mathbb{R}^d \times \mathbb{R}^d} f \, dv \, dx = m$, then it coincides with the critical point $f = \mathcal{M}_{\text{eq}}$, $\Psi = \Psi_{\text{eq}}$ and $\pi = 0$.*

It follows from this, together with the uniqueness of the critical point, that there is at most one minimizer. That such a minimizer exists and that, therefore, the unique critical point is indeed a minimizer, will be proven below, in Lemma 3.4.

Proof. By virtue of Lemma 2.1 \mathcal{J} is bounded from below, when working on non negative functions f with fixed total mass $m > 0$ (see the manipulation in Appendix A). Let (f, Ψ, π) be a minimizer of \mathcal{J} with the constraint $\iint_{\mathbb{R}^d \times \mathbb{R}^d} f \, dv \, dx = m$. Since $\mathcal{J}(f, \Psi, \pi)$ is finite, we deduce that $f(x, v) \geq 0$ a. e. and $h(f) \in L^1(\mathbb{R}^d \times \mathbb{R}^d)$. We readily check that $\pi = 0$ and $-\Delta_z \Psi(x, z) = -\sigma_2(z) \left(\sigma_1 \star \int_{\mathbb{R}^d} f(\cdot, v) \, dv \right)(x)$, just by expanding $\mathcal{J}(f, \Psi + \delta\psi, \pi) - \mathcal{J}(f, \Psi, \pi) \geq 0$ and $\mathcal{J}(f, \Psi, \pi + \delta\tilde{\pi}) - \mathcal{J}(f, \Psi, \pi) \geq 0$ with $\psi, \tilde{\pi} \in C_c^\infty(\mathbb{R}^d \times \mathbb{R}^n)$, and letting δ go to 0. It remains to find the expression of the distribution function f . We follow the arguments detailed in [19, 24]. We start by discussing the case where h is a C^1 function over $[0, \infty)$. For $\epsilon > 0$, we set

$$S_\epsilon = \{(x, v) \in \mathbb{R}^d \times \mathbb{R}^d, \epsilon \leq f(x, v) \leq 1/\epsilon\}.$$

We consider $w \in L^\infty(\mathbb{R}^d \times \mathbb{R}^d)$, with $\text{supp}(w) \subset B(0, R)$, for some $0 < R < \infty$ and such that $w \geq 0$ on $\mathbb{C}S_\epsilon$. As a consequence of the mass constraint, and the Bienaymé-Tchebychev inequality, we have, for small enough ϵ 's,

$$0 < \text{meas}(S_\epsilon) \leq \frac{m}{\epsilon} < \infty.$$

Then, we can find $\tau(\epsilon) > 0$ such that, for any $0 < \tau \leq \tau(\epsilon)$ the function

$$f_\tau(x, v) = f(x, v) + \tau \left(w(x, v) - \mathbf{1}_{S_\epsilon}(x, v) \frac{\int_{\mathbb{R}^d \times \mathbb{R}^d} w(y, \xi) \, dy \, d\xi}{\text{meas}(S_\epsilon)} \right)$$

is non-negative and satisfies $\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_\tau \, dv \, dx = m$. Accordingly we have

$$\mathcal{J}(f_\tau, \Psi, \pi) - \mathcal{J}(f, \Psi, \pi) \geq 0.$$

Since h is C^1 and $f_\tau - f$ is supported on a set of finite measure, it follows

that

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(h'(f(x, v)) + \frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \times \left(w(x, v) - \mathbf{1}_{S_\epsilon}(x, v) \frac{\int_{\mathbb{R}^d \times \mathbb{R}^d} w(y, \xi) \, dy \, d\xi}{\text{meas}(S_\epsilon)} \right) \, dv \, dx \geq 0$$

which recasts as

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} w(x, v) \left(h'(f(x, v)) + \frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) + \lambda_\epsilon \right) \, dv \, dx \geq 0$$

where we have set

$$\lambda_\epsilon = -\frac{1}{\text{meas}(S_\epsilon)} \int_{S_\epsilon} \left(h'(f(y, w)) + \frac{|w|^2}{2} + V(y) + \Phi[\Psi](y) \right) \, dy \, dw \in \mathbb{R}.$$

Since this relation holds for any trial function w verifying the conditions prescribed above, we deduce the following properties:

- On the one hand, for a. e. $(x, v) \in S_\epsilon$, we have

$$h'(f(x, v)) + \frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) = -\lambda_\epsilon$$

which, additionally, tells us that $\lambda_\epsilon = \lambda$ does not depend on ϵ .

- On the other hand, on $\complement S_\epsilon$, we get $h'(f(x, v)) + \frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) + \lambda \geq 0$.

Letting ϵ go to 0, we conclude that either $f(x, v) > 0$, and $h'(f(x, v)) = -\frac{|v|^2}{2} - V(x) - \Phi[\Psi](x) - \lambda > \eta_0$, or $f(x, v) = 0$ and $\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) + \lambda \geq -\eta_0$. We recover the formula for the critical points of \mathcal{J} .

The argument needs to be slightly adapted to handle the case $h(s) = s \ln(s) - s$, for which $h'(0)$ is not defined. We therefore need to exclude the possibility that $f(x, v) = 0$. The proof therefore begins with the justification that f takes strictly positive values only. We argue by contradiction: suppose that the set $\mathcal{O} = \{(x, v) \in \mathbb{R}^d \times \mathbb{R}^d, f(x, v) = 0\}$ has a positive measure (possibly $+\infty$). The regularity of the Lebesgue measure allows us to find a compact set $K \subset \mathcal{O}$ such that $\text{meas}(K) > 0$. We now set for $\tau > 0$

$$f_\tau(x, v) = f(x, v) + \tau \left(\mathbf{1}_K - \mathbf{1}_{S_\epsilon} \frac{\text{meas}(K)}{\text{meas}(S_\epsilon)} \right),$$

where S_ϵ is defined as above. Clearly $K \cap S_\epsilon = \emptyset$ and, again, we can find $\tau(\epsilon) > 0$ such that f_τ remains non negative for any $0 < \tau \leq \tau(\epsilon)$, while $\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_\tau \, dv \, dx = m$. Therefore, we have

$$\begin{aligned} 0 &\leq \mathcal{J}(f_\tau, \Psi, \pi) - \mathcal{J}(f, \Psi, \pi) \\ &\leq \iint_{\mathbb{R}^d \times \mathbb{R}^d} (h(f_\tau) - h(f)) \, dv \, dx \\ &\quad + \tau \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \left(\mathbf{1}_K - \mathbf{1}_{S_\epsilon} \frac{\text{meas}(K)}{\text{meas}(S_\epsilon)} \right) \, dv \, dx. \end{aligned}$$

By definition $f_\tau(x, v) = f(x, v)$ on $\mathcal{C}(K \cup S_\epsilon)$, $f_\tau(x, v) = f(x, v) + \tau = \tau$ on K and $f_\tau(x, v) = f(x, v) - \tau \frac{\text{meas}(K)}{\text{meas}(S_\epsilon)}$ on S_ϵ . This yields

$$\begin{aligned} 0 &\leq \text{meas}(K)(\tau \ln(\tau) - \tau) + \tau \iint_K \left(\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \, dv \, dx \\ &\quad + \iint_{S_\epsilon} (h(f_\tau) - h(f)) \, dv \, dx \\ &\quad - \tau \frac{\text{meas}(K)}{\text{meas}(S_\epsilon)} \iint_{S_\epsilon} \left(\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \, dv \, dx. \end{aligned}$$

Let us use a shorthand notation for the real number

$$\begin{aligned} A &= \iint_K \left(\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \, dv \, dx \\ &\quad - \frac{\text{meas}(K)}{\text{meas}(S_\epsilon)} \iint_{S_\epsilon} \left(\frac{|v|^2}{2} + V(x) + \Phi[\Psi](x) \right) \, dv \, dx. \end{aligned}$$

Dividing the previous relation by $\tau > 0$, we arrive at

$$0 \leq \text{meas}(K)(\ln(\tau) - 1) + A + \iint_{S_\epsilon} \frac{h(f_\tau) - h(f)}{\tau} \, dv \, dx.$$

As τ tends to 0 the last term tends to $\iint_{S_\epsilon} \ln(f) \, dv \, dx$, which is also finite for any fixed $\epsilon > 0$. We are led to a contradiction since $\lim_{\tau \rightarrow 0} \ln(\tau) = -\infty$. We conclude that $\text{meas}(\mathcal{O}) = 0$. From here, we can repeat the previous argument to identify f by means of the Lagrange multiplier and to recover this way the formula that defines the critical point. \blacksquare

3. Dynamical stability

To start, it is convenient to make the relative entropy with respect to the equilibrium state appear. For h a given strictly convex function, we set

$$H(f|g) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} (h(f) - h(g) - h'(g)(f - g)) \, dv \, dx.$$

This quantity measures how far f is from g since we have

$$H(f|g) \geq 0 \text{ and } H(f|g) = 0 \text{ iff } f = g.$$

Remark 3.1. *It is worth pointing out that, for the case of $h(s) = s \ln(s) - s$, the functional H controls the L^1 norm of $f - g$, by virtue of the Csiszar-Kullback inequality [7, 20]. For other entropies h , it is possible to deduce L^p estimates, see [6].*

In the specific case where $g = \mathcal{M}_{\text{eq}}$, we can use

$$h'(\mathcal{M}_{\text{eq}}(x, v)) = -\frac{|v|^2}{2} - V(x) - \Phi_{\text{eq}}(x) - \lambda_{\text{eq}}$$

and we arrive at

$$H(f|\mathcal{M}_{\text{eq}}) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(h(f) - h(\mathcal{M}_{\text{eq}}) + (f - \mathcal{M}_{\text{eq}}) \left(\frac{|v|^2}{2} + V(x) + \Phi_{\text{eq}}(x) \right) \right) dv dx$$

provided

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f dv dx = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \mathcal{M}_{\text{eq}} dv dx = m.$$

We now compute

$$\mathcal{J}(f, \Psi, \pi) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) = I + II + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi|^2 dz dx$$

with, on the one hand

$$\begin{aligned} I &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(h(f) + \left(\frac{|v|^2}{2} + V(x) \right) f - h(\mathcal{M}_{\text{eq}}) - \left(\frac{|v|^2}{2} + V(x) \right) \mathcal{M}_{\text{eq}} \right) dv dx \\ &= H(f|\mathcal{M}_{\text{eq}}) + \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\mathcal{M}_{\text{eq}} - f) \Phi_{\text{eq}} dv dx \end{aligned}$$

and, on the other hand

$$\begin{aligned}
II &= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(\Phi[\Psi]f - \Phi_{\text{eq}} \mathcal{M}_{\text{eq}} \right) dv dx \\
&\quad + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} \left(|\nabla_z \Psi|^2 - |\nabla_z \Psi_{\text{eq}}|^2 \right) dz dx \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi_{\text{eq}}(f - \mathcal{M}_{\text{eq}}) dv dx + \iint_{\mathbb{R}^d \times \mathbb{R}^d} f(\Phi[\Psi] - \Phi_{\text{eq}}) dv dx \\
&\quad + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx \\
&\quad - c^2 \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z \Psi_{\text{eq}}|^2 dz dx + c^2 \iint_{\mathbb{R}^d \times \mathbb{R}^n} \nabla_z \Psi \cdot \nabla_z \Psi_{\text{eq}} dz dx \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi_{\text{eq}}(f - \mathcal{M}_{\text{eq}}) dv dx \\
&\quad + \iint_{\mathbb{R}^d \times \mathbb{R}^d} f \sigma_1 \star \left(\int_{\mathbb{R}^n} \sigma_2(\Psi - \Psi_{\text{eq}}) dz \right) dv dx \\
&\quad + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx \\
&\quad + c^2 \iint_{\mathbb{R}^d \times \mathbb{R}^n} (\Psi_{\text{eq}} - \Psi) \cdot \Delta_y \Psi_{\text{eq}} dz dx \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi_{\text{eq}}(f - \mathcal{M}_{\text{eq}}) dv dx \\
&\quad + \iint_{\mathbb{R}^d \times \mathbb{R}^d} f \sigma_1 \star \left(\int_{\mathbb{R}^n} \sigma_2(\Psi - \Psi_{\text{eq}}) dz \right) dv dx \\
&\quad + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx \\
&\quad + c^2 \iint_{\mathbb{R}^d \times \mathbb{R}^n} (\Psi_{\text{eq}} - \Psi) \sigma_2 \sigma_1 \star \left(\int_{\mathbb{R}^d} \mathcal{M}_{\text{eq}} dv \right) dz dx \\
&= \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi_{\text{eq}}(f - \mathcal{M}_{\text{eq}}) dv dx + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx \\
&\quad + \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(\Psi - \Psi_{\text{eq}}) \sigma_1 \star (f - \mathcal{M}_{\text{eq}}) dv dz dx.
\end{aligned}$$

We end up with

$$\begin{aligned}
& \mathcal{J}(f, \Psi, \pi) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) \\
&= H(f|\mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi|^2 dz dx \\
&\quad + \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(z) (\Psi - \Psi_{\text{eq}})(x, z) \sigma_1 \star (f - \mathcal{M}_{\text{eq}})(x, v) dv dz dx.
\end{aligned} \tag{13}$$

The first three terms in the right hand side are non negative; only the last integral, which is nothing but

$$\mathcal{B}\left(\Psi - \Psi_{\text{eq}}, \int_{\mathbb{R}^d} f - \mathcal{M}_{\text{eq}} dv\right),$$

does not have a definite sign. The main result of this paper can be stated as follows.

Theorem 3.2. *Let $m > 0$ and suppose **(H0)**–**(H3)**. Let $c > c_0$, as in Proposition 2.3. For any $\epsilon > 0$, there exists $\eta > 0$ such that, if the initial data*

$$f|_{t=0} = f_0, \quad \Psi|_{t=0} = \Psi_0, \quad \partial_t \Psi|_{t=0} = \Psi_1$$

for (1a)–(1d) satisfies

$$H(f_0|\mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi_0 - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\Psi_1|^2 dz dx \leq \eta$$

then, for any $t \geq 0$, we have

$$\begin{aligned}
& H(f(t, \cdot)|\mathcal{M}_{\text{eq}}) \\
&+ \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi(t, x, z) - \Psi_{\text{eq}}(x, z))|^2 dz dx \\
&+ \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\partial_t \Psi(t, x, z)|^2 dz dx \leq \epsilon.
\end{aligned}$$

Remark 3.3. *Let us make a couple of comments:*

- (a) *The preliminary result in [2, Appendix A] is an analogous statement to the one in Theorem 3.2, for the linearized equation and the equilibrium (3); in that case, a weighted L^2 norm on the particle distribution is used, which corresponds to the linearization of an appropriate relative entropy H .*

(b) As explained in [5], the condition that the propagation speed c of the waves must be large can be understood to mean that the environment can evacuate energy quickly to infinity. Whether this condition is necessary for the stability result above is not clear. It is not needed in [5] to prove stability of a single particle coupled to a vibrating environment in the presence of a confining potential. But, as pointed out before, the situation here is more complex. Even the existence–uniqueness of the equilibria we study here is guaranteed only for large enough c (see Proposition 2.3 and [2, Theorem 2.1] for the issue of the uniqueness of the equilibria with prescribed mass). Here, the equilibria are defined as ground states of the energy–Casimir functionals, which gives a correspondence between the shape of the equilibrium and the functional. The next ingredient in the proof of their existence and uniqueness relies on compactness properties of minimizing sequences. This approach thus implies some restrictions on the class of considered equilibria; we refer the reader to [22] for a thorough introduction to the stability issues for the Vlasov–Poisson system and possible generalizations.

The main argument for proving Theorem 3.2 is the following characterization of the minimizers of \mathcal{J} , which completes Lemma 2.4.

Lemma 3.4 (Minimization of \mathcal{J}). *The following assertions hold:*

- (i) *The functional \mathcal{J} has a unique minimizer over (12) with a particle distribution function with mass m and it is $(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}})$.*
- (ii) *Let $((g^\nu, \Psi^\nu, \pi^\nu))_{\nu \in \mathbb{N}}$ be a minimizing sequence of the functional \mathcal{J} , among the distributions with mass m ; then, we can extract a subsequence $(\nu_k)_{k \in \mathbb{N}}$ such that $(g^{\nu_k}, \Psi^{\nu_k}, \pi^{\nu_k})$ converges weakly to the minimizer $(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}})$ and we have*

$$\lim_{k \rightarrow \infty} \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(z) (\Psi^{\nu_k} - \Psi_{\text{eq}})(x, z) \times \sigma_1 \star (g^{\nu_k} - \mathcal{M}_{\text{eq}})(x, v) \, dv \, dz \, dx = 0.$$

Proof. We have seen that $(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}})$ is the unique critical point of \mathcal{J} among the distributions with mass m (see Proposition 2.3). By Lemma 2.4, if \mathcal{J} has a minimizer it has to be $(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}})$. In order to prove (i), we just have to prove that \mathcal{J} admits at least one minimizer.

To this end, let us consider $((g^\nu, \Psi^\nu, \pi^\nu))_{\nu \in \mathbb{N}}$, a minimizing sequence of \mathcal{J} . As explained in Appendix A, it follows from the boundedness of the functional \mathcal{J} , together with the fact that the coupling term is bounded (see Lemma 2.1), that we can suppose, for a suitable subsequence,

- $g^{\nu_k} \rightharpoonup g$ weakly in $L^1(\mathbb{R}^d \times \mathbb{R}^d)$,
- $\nabla_z \Psi^{\nu_k} \rightharpoonup \nabla_z \Psi$ weakly in $L^2(\mathbb{R}^d \times \mathbb{R}^n)$,
- $\pi^{\nu_k} \rightharpoonup \pi$ weakly in $L^2(\mathbb{R}^d \times \mathbb{R}^n)$.

In fact, the weak compactness of $(g^\nu)_{\nu \in \mathbb{N}}$ in L^1 follows from an application of the Dunford–Pettis theorem, see [16, Section 7.3.2], by using **(H0)** and **(H1)**. This is direct when h is non negative; for $h(s) = s \ln(s) - s$, the necessary estimate relies on a standard trick which is detailed in Appendix A.

We are going to prove that the limit (g, Ψ, π) minimizes \mathcal{J} . It is convenient to work with the expression of \mathcal{J} in (13). Indeed, owing to convexity properties (see for instance [4, Corollary III.8]), we already get

$$\begin{aligned} & \liminf_{k \rightarrow \infty} \left\{ H(g^{\nu_k} | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi^{\nu_k} - \Psi_{\text{eq}})|^2 dz dx \right. \\ & \qquad \qquad \qquad \left. + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi_{\nu_k}|^2 dz dx \right\} \\ & \geq H(g | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi|^2 dz dx. \end{aligned} \tag{14}$$

We also have, as $k \rightarrow \infty$,

$$\begin{aligned} \rho^{\nu_k}(x) &= \int_{\mathbb{R}^d} g^{\nu_k}(x, v) dv \rightharpoonup \rho(x) = \int_{\mathbb{R}^d} g(x, v) dv \text{ weakly in } L^1(\mathbb{R}^d), \\ \iint_{\mathbb{R}^d \times \mathbb{R}^d} g^{\nu_k}(x, v) dv dx &= m = \iint_{\mathbb{R}^d \times \mathbb{R}^d} g(x, v) dv dx. \end{aligned}$$

Accordingly, we observe that, for any $\varphi \in L^\infty(\mathbb{R}^d)$,

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{\mathbb{R}^d} \sigma_1 \star \rho^{\nu_k} \varphi dx &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}^d} \rho^{\nu_k} \sigma_1 \star \varphi dx \\ &= \int_{\mathbb{R}^d} \rho \sigma_1 \star \varphi dx = \int_{\mathbb{R}^d} \sigma_1 \star \rho \varphi dx, \end{aligned}$$

since, with $\sigma_1 \in L^1 \cap L^\infty(\mathbb{R}^d)$, $\sigma_1 \star \varphi$ belongs to $L^\infty(\mathbb{R}^d)$. It means that $\sigma_1 \star \rho^{\nu_k}$ converges weakly to $\sigma_1 \star \rho$ in $L^1(\mathbb{R}^d)$. Furthermore, since for a. e. $x \in \mathbb{R}^d$, the function $y \mapsto \sigma_1(x - y)$ lies in $L^\infty(\mathbb{R}^d)$, the convergence $\sigma_1 \star \rho^{\nu_k}(x) \rightarrow \sigma_1 \star \rho(x)$ also holds for a. e. $x \in \mathbb{R}^d$. Combining the two informations we deduce that $\sigma_1 \star \rho^{\nu_k}$ converges strongly to $\sigma_1 \star \rho$ in $L^1(\mathbb{R}^d)$ see [16, Theorem 7.60]. Since $\sigma_1 \in L^1 \cap L^\infty(\mathbb{R}^d)$, $\sigma_1 \star \rho^{\nu_k}$ is actually bounded in any $L^p(\mathbb{R}^d)$ spaces, $1 \leq p \leq \infty$, and the convergence holds strongly in $L^p(\mathbb{R}^d)$

for any finite p . We turn now to the behavior of $\int_{\mathbb{R}^n} \sigma_2(z) \Psi^{\nu_k}(x, z) dz$. The Gagliardo-Nirenberg-Sobolev inequality, that we already used for proving Lemma 2.1, tells us that Ψ^{ν_k} is bounded in $L^2(\mathbb{R}^d; L^{2n/(n-2)}(\mathbb{R}^n))$. Hence, applying [15, Theorem 8.20.5] and the Banach-Alaoglu-Bourbaki theorem, we can assume that it tends to ψ , weakly in this space. Therefore, for any $\varphi \in L^2(\mathbb{R}^d)$, the product $\sigma_2(z)\varphi(x)$ lies in $L^2(\mathbb{R}^d; L^{2n/(n+2)}(\mathbb{R}^n))$ and we get

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{\mathbb{R}^d} \varphi(x) \left(\int_{\mathbb{R}^n} \sigma_2(z) \Psi^{\nu_k}(x, z) dz \right) dx \\ = \int_{\mathbb{R}^d} \varphi(x) \left(\int_{\mathbb{R}^n} \sigma_2(z) \Psi(x, z) dz \right) dx. \end{aligned}$$

In other words, $\int_{\mathbb{R}^n} \sigma_2(z) \Psi^{\nu_k}(x, z) dz$ converges to $\int_{\mathbb{R}^n} \sigma_2(z) \Psi(x, z) dz$ weakly in $L^2(\mathbb{R}^d)$. We conclude that

$$\begin{aligned} \lim_{k \rightarrow \infty} \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(\Psi^{\nu_k} - \Psi_{\text{eq}})(x, z) \sigma_1 \star (g^{\nu_k} - \mathcal{M}_{\text{eq}})(x, v) dv dz dx \\ = \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(\Psi - \Psi_{\text{eq}})(x, z) \sigma_1 \star (g - \mathcal{M}_{\text{eq}})(x, v) dv dz dx. \end{aligned} \quad (15)$$

We combine this relation to (14) and going back to (13) we deduce that

$$\liminf_{k \rightarrow \infty} \mathcal{J}(g^{\nu_k}, \Psi^{\nu_k}, \pi^{\nu_k}) \geq \mathcal{J}(g, \Psi, \pi).$$

Since $((g^\nu, \Psi^\nu, \pi^\nu))_{\nu \in \mathbb{N}}$ is a minimizing sequence of \mathcal{J} , we deduce that (g, Ψ, π) is a minimizer of \mathcal{J} , with g verifying $\iint_{\mathbb{R}^d \times \mathbb{R}^d} g dv dx = m$. This already proves (i). As said above, according to Proposition 2.3 and Lemma 2.4, we get

$$g = \mathcal{M}_{\text{eq}}, \quad \psi = \Psi_{\text{eq}}, \quad \pi = \pi_{\text{eq}} = 0.$$

Going back to (15), we obtain

$$\lim_{k \rightarrow \infty} \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(z) (\Psi^{\nu_k} - \Psi_{\text{eq}})(x, z) \sigma_1 \star (g^{\nu_k} - \mathcal{M}_{\text{eq}})(x, v) dv dz dx = 0$$

It ends the proof of (ii). \blacksquare

Proof of Theorem 3.2. We argue by contradiction. We thus assume that we can find:

- a sequence of initial data $((f_0^\nu, \Psi_0^\nu, \Psi_1^\nu))_{\nu \in \mathbb{N}}$ such that

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} f_0^\nu dv dx = m$$

and

$$\lim_{\nu \rightarrow \infty} \left\{ H(f_0^\nu | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi_0^\nu - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\Psi_1^\nu|^2 dz dx \right\} = 0, \quad (16)$$

- a positive number $\epsilon > 0$ and a sequence of times $(t^\nu)_{\nu \in \mathbb{N}}$

such that $g^\nu(x, v) = f^\nu(t^\nu, x, v)$, $\tilde{\Psi}^\nu(x, z) = \Psi^\nu(t^\nu, x, z)$, and $\pi^\nu(x, z) = \partial_t \Psi^\nu(t^\nu, x, z)$, with (f^ν, Ψ^ν) the solution of (1a)–(1d) determined by

$$f^\nu|_{t=0} = f_0^\nu, \quad \Psi^\nu|_{t=0} = \Psi_0^\nu, \quad \partial_t \Psi^\nu|_{t=0} = \Psi_1^\nu,$$

satisfy

$$H(g^\nu | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\tilde{\Psi}^\nu - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi^\nu|^2 dz dx \geq \epsilon. \quad (17)$$

Conservation of the total energy (7) and conservation of the Casimir functionals (9) imply that

$$\mathcal{J}(g^\nu, \tilde{\Psi}^\nu, \pi^\nu) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) = \mathcal{J}(f_0^\nu, \Psi_0^\nu, \Psi_1^\nu) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) \quad (18)$$

and, of course, by mass conservation, we have

$$\iint_{\mathbb{R}^d \times \mathbb{R}^d} g^\nu dv dx = m. \quad (19)$$

Using (13), relation (18) can be rewritten as

$$\begin{aligned} & H(g^\nu | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\tilde{\Psi}^\nu - \Psi_{\text{eq}})|^2 dz dx + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi^\nu|^2 dz dx \\ & + \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(\tilde{\Psi}^\nu - \Psi_{\text{eq}}) \sigma_1 \star (g^\nu - \mathcal{M}_{\text{eq}}) dv dz dx \\ & = H(f_0^\nu | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z(\Psi_0^\nu - \Psi_{\text{eq}})|^2 dz dx \\ & + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\Psi_1^\nu|^2 dz dx \\ & + \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(z) (\Psi_0^\nu - \Psi_{\text{eq}})(x, z) \\ & \quad \times \sigma_1 \star (f_0^\nu - \mathcal{M}_{\text{eq}})(x, v) dv dz dx. \end{aligned} \quad (20)$$

We start by observing that the right hand side tends to 0 as $\nu \rightarrow \infty$.

Lemma 3.5. *We have*

$$\lim_{\nu \rightarrow \infty} \mathcal{J}(f_0^\nu, \Psi_0^\nu, \Psi_1^\nu) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) = 0.$$

Proof. This is a consequence of (16). The last term in (20) is nothing but $\mathcal{B}(\Psi_0^\nu - \Psi_{\text{eq}}, \rho_0^\nu - \rho_{\text{eq}})$, with $\rho_0^\nu(x) = \int_{\mathbb{R}^d} f_0^\nu(x, v) dv$, and we appeal to the continuity property stated in Lemma 2.1. ■

It follows that

$$\lim_{\nu \rightarrow \infty} \mathcal{J}(g^\nu, \tilde{\Psi}^\nu, \pi^\nu) = \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) \quad (21)$$

We are led to a contradiction owing to Lemma 3.4 which makes $((g^\nu, \tilde{\Psi}^\nu, \pi^\nu))_{\nu \in \mathbb{N}}$ appear as a minimizing sequence for \mathcal{J} with the mass constrained to be m . By using Lemma 3.4-(ii), we can find a suitable subsequence $(\nu_k)_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} \iiint_{\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^n} \sigma_2(z) (\tilde{\Psi}^{\nu_k} - \Psi_{\text{eq}})(x, z) \sigma_1 \star (g^{\nu_k} - \mathcal{M}_{\text{eq}})(x, v) dv dz dx = 0.$$

Going back to (18) and (20), we are thus led to

$$\begin{aligned} & \lim_{k \rightarrow \infty} \left\{ \mathcal{J}(g^{\nu_k}, \tilde{\Psi}^{\nu_k}, \pi^{\nu_k}) - \mathcal{J}(\mathcal{M}_{\text{eq}}, \Psi_{\text{eq}}, \pi_{\text{eq}}) \right\} = 0 \\ & = \lim_{k \rightarrow \infty} \left\{ H(g^{\nu_k} | \mathcal{M}_{\text{eq}}) + \frac{c^2}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\nabla_z (\tilde{\Psi}^{\nu_k} - \Psi_{\text{eq}})|^2 dz dx \right. \\ & \quad \left. + \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^n} |\pi^{\nu_k}|^2 dz dx \right\}. \end{aligned}$$

Since the three sequences involved in the last expression are all non negative, we conclude that

$$\begin{aligned} H(g^{\nu_k} | \mathcal{M}_{\text{eq}}) & \xrightarrow[k \rightarrow \infty]{} 0, \\ \|\nabla_z (\tilde{\Psi}^{\nu_k} - \Psi_{\text{eq}})\|_{L^2(\mathbb{R}^d \times \mathbb{R}^n)} & \xrightarrow[k \rightarrow \infty]{} 0, \quad \|\pi^{\nu_k}\|_{L^2(\mathbb{R}^d \times \mathbb{R}^n)} \xrightarrow[k \rightarrow \infty]{} 0. \end{aligned}$$

This contradicts (17). ■

Appendix A. Weak compactness in L^1 for $h(s) = s \ln(s) - s$

The difficulty comes from the fact that $h(f)$ might change sign. We shall prove that $g^\nu |\ln(g^\nu)|$ and $(|v|^2 + V(x))g^\nu$ are bounded in $L^1(\mathbb{R}^d \times \mathbb{R}^d)$, which will allow us to apply the Dunford–Pettis theorem [16, Section 7.3.2], taking into account **(H0)**.

Owing to Young's inequality, we deduce from Lemma 2.1 that there exists $C > 0$ such that for any $\beta > 0$, we have $\mathcal{B}(g^\nu, \psi^\nu) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \Phi[\psi^\nu] g^\nu \, dv \, dx \geq -\beta \|\nabla \psi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 - \frac{C}{\beta} m^2$. Therefore we get

$$\begin{aligned} & \frac{C}{\beta} m^2 + \mathcal{J}(g^\nu, \psi^\nu, \pi^\nu) \\ & \geq \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(g^\nu) \, dv \, dx + \iint_{\mathbb{R}^d \times \mathbb{R}^d} g^\nu \left(\frac{|v|^2}{2} + V(x) \right) \, dv \, dx \\ & \quad + \left(\frac{c^2}{2} - \beta \right) \|\nabla \psi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 + \frac{1}{2} \|\pi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2, \end{aligned}$$

where we can pick $0 < \beta < c^2/2$. Since the left hand side is bounded with respect to ν , it already justifies the weak compactness of g^ν , ψ^ν and π^ν when h is non negative and satisfies **(H1)**.

For $h(s) = s \ln(s) - s$, we use the following argument: for $\Omega \geq 0$, we have

$$\begin{aligned} s |\ln(s)| &= s \ln(s) - 2s \ln(s) (\mathbf{1}_{e^{-\Omega} \leq s \leq 1} + \mathbf{1}_{e^{-\Omega} > s}) \\ &\leq s \ln(s) + 2\Omega s + \frac{4}{e} e^{-\Omega/2}. \end{aligned}$$

We use this inequality with $\Omega = 2\alpha \left(\frac{|v|^2}{2} + V(x) \right)$, $0 < \alpha < 1/4$. It allows us to obtain

$$\begin{aligned} & \iint_{\mathbb{R}^d \times \mathbb{R}^d} g^\nu |\ln(g^\nu)| \, dv \, dx + (1 - 4\alpha) \iint_{\mathbb{R}^d \times \mathbb{R}^d} g^\nu \left(\frac{|v|^2}{2} + V(x) \right) \, dv \, dx \\ & \quad + \left(\frac{c^2}{2} - \beta \right) \|\nabla \psi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 + \frac{1}{2} \|\pi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 \\ & \leq \iint_{\mathbb{R}^d \times \mathbb{R}^d} h(g^\nu) \, dv \, dx + \iint_{\mathbb{R}^d \times \mathbb{R}^d} g^\nu \left(\frac{|v|^2}{2} + V(x) \right) \, dv \, dx \\ & \quad + \frac{c^2}{2} \|\nabla \psi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 + \frac{1}{2} \|\pi^\nu\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)}^2 + \mathcal{B}(g^\nu, \psi^\nu) \\ & \quad + m + \frac{C}{\beta} m^2 + \frac{4}{e} \iint_{\mathbb{R}^d \times \mathbb{R}^d} e^{-\alpha V(x)} e^{-\alpha |v|^2/2} \, dv \, dx \\ & \leq \mathcal{J}(g^\nu, \psi^\nu, \pi^\nu) + m + \frac{C}{\beta} m^2 + \frac{4}{e} \iint_{\mathbb{R}^d \times \mathbb{R}^d} e^{-\alpha V(x)} e^{-\alpha |v|^2/2} \, dv \, dx \end{aligned}$$

which is bounded uniformly with respect to ν .

- [1] Aguer, B., De Bièvre, S., Lafitte, P., Parris, P. E., 2010. Classical motion in force fields with short range correlations. *J. Stat. Phys.* 138 (4-5), 780–814.

- [2] Alonso, R., Goudon, T., Vasseur, A., 2017. Damping of particles interacting with a vibrating medium. *Ann. IHP. Anal. Non-Linéaire* 34 (7), 1727–1758.
- [3] Ben Abdallah, N., Dolbeault, J., 2003. Relative entropies for kinetic equations in bounded domains (irreversibility, stationary solutions, uniqueness). *Archiv. Ration. Mech. Anal.* 168 (4), 253–298.
- [4] Brézis, H., 1987. *Analyse fonctionnelle. Théorie et applications.* Masson.
- [5] Bruneau, L., De Bièvre, S., 2002. A Hamiltonian model for linear friction in a homogeneous medium. *Comm. Math. Phys.* 229 (3), 511–542.
- [6] Cáceres, M.-J., Carrillo, J. A., Dolbeault, J., 2002. Nonlinear stability in L^p for a confined system of charged particles. *SIAM J. Math. Anal.* 34, 478–494.
- [7] Csiszar, I., 1967. Information-type measures of difference of probability distributions and indirect observations. *Studia Sci. Math. Hungar.* 2, 299–318.
- [8] De Bièvre, S., Goudon, T., Vasseur, A., 2016. Particles interacting with a vibrating medium: existence of solutions and convergence to the Vlasov–Poisson system. *SIAM J. Math. Anal.* 48 (6), 3984–4020.
- [9] De Bièvre, S., Lafitte, P., Parris, P. E., 2007. Normal transport at positive temperatures in classical Hamiltonian open systems. In: *Adventures in mathematical physics.* Vol. 447 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, pp. 57–71.
- [10] De Bièvre, S., Parris, P. E., 2011. Equilibration, generalized equipartition, and diffusion in dynamical Lorentz gases. *J. Stat. Phys.* 142 (2), 356–385.
- [11] De Bièvre, S., Parris, P. E., Silviu, A., 2005. Chaotic dynamics of a free particle interacting linearly with a harmonic oscillator. *Phys. D* 208 (1-2), 96–114.
- [12] Dolbeault, J., Markowich, P., Unterreiter, A., 2001. On singular limits of mean-field equations. *Arch. Rational Mech. Anal.* 158 (4), 319–351.
- [13] Dolbeault, J., Mouhot, C., Schmeiser, C., 2015. Hypocoercivity for linear kinetic equations conserving mass. *Trans. AMS* 367 (6), 3807–3828.

- [14] Dolbeault, J., Sánchez, O., Soler, J., 2004. Asymptotic behaviour for the Vlasov–Poisson system in the stellar-dynamics case. *Archiv. Ration. Mech. Anal.* 171, 301–327.
- [15] Edwards, R. E., 1995. *Functional analysis*. Dover Publications Inc., New York, theory and applications, Corrected reprint of the 1965 original.
- [16] Goudon, T., 2011. *Intégration: Intégrale de Lebesgue et introduction à l’analyse fonctionnelle*. Références Sciences. Ellipses.
- [17] Goudon, T., Vavasseur, A., 2016. Mean field limit for particles interacting with a vibrating medium. *Annali Univ. Ferrara* 62 (2), 231–273.
- [18] Guo, Y., 1999. Variational method for stable polytropic galaxies. *Archiv. Ration. Mech. Anal.* 130, 163–182.
- [19] Guo, Y., Rein, G., 1999. Stable steady states in stellar dynamics. *Archiv. Ration. Mech. Anal.* 147, 225–243.
- [20] Kullback, S., 1967. A lower bound for discrimination information in terms of variation. *IEEE Trans. Information Theory* 4, 126–127.
- [21] Lafitte, P., Parris, P. E., De Bièvre, S., 2008. Normal transport properties in a metastable stationary state for a classical particle coupled to a non-Ohmic bath. *J. Stat. Phys.* 132 (5), 863–879.
- [22] Mouhot, C., 2011-2012. Stabilité orbitale pour le système de Vlasov-Poisson gravitationnel. Séminaire Bourbaki, Exposé 1044.
- [23] Nirenberg, L., 1959. On elliptic partial differential equations. *Ann. Scuola Norm. Sup. Pisa, Classe di Scienze* 13 (2), 115–162.
- [24] Rein, G., 2007. Collisionless kinetic equations from astrophysics: The Vlasov–Poisson system. In: *Handbook of Differential Equations: Evolutionary Equations*. Vol. 3. Elsevier, pp. 385–476.
- [25] Soret, E., De Bièvre, S., 2015. Stochastic acceleration in a random time-dependent potential. *Stochastic Process. Appl.* 125 (7), 2752–2785.
URL <http://dx.doi.org/10.1016/j.spa.2015.01.012>
- [26] Vavasseur, A., 2016. Some models of particles interacting with their environment. Ph.D. thesis, University Nice Sophia Antipolis.
- [27] Wolansky, G., 1999. On nonlinear stability of polytropic galaxies. *Ann. Inst. H. Poincaré* 16, 15–48.