

Derivation Graphs, Greediness, and Bounded-treewidth in the Context of Existential Rules

Tim Lyon

(Joint Work with Sebastian Rudolph)

Computational Logic Group, Institute of Artificial Intelligence,
Technische Universität Dresden



- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion

- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z(\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z(\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z(\text{female}(z) \wedge \text{parent}(z, x))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(\textcolor{red}{x}) \wedge \text{human}(\textcolor{red}{x}) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, \textcolor{red}{x}))$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{ \text{male}(\text{Joe}), \text{human}(\text{Joe}) \}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{ \text{male}(\text{Joe}), \text{human}(\text{Joe}) \}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, \textcolor{red}{h}_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = male(x) \wedge human(x) \rightarrow \exists z (female(z) \wedge parent(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{male(Joe), human(Joe)\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Example: Let $h(x) = \text{Joe}$.

$$1 \quad F_0 = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Example: Let $h(x) = \text{Joe}$.

$$1 \quad F_0 = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

$$2 \quad F_1 = \{\text{male}(\text{Joe}), \text{human}(\text{Joe}), \text{female}(z), \text{parent}(z, \text{Joe})\}$$

Preliminaries

Existential Rule: $\forall \vec{x} \forall \vec{y} (\phi(\vec{x}, \vec{y}) \rightarrow \exists \vec{z} \psi(\vec{y}, \vec{z}))$

Example:

$$\rho = \text{male}(x) \wedge \text{human}(x) \rightarrow \exists z (\text{female}(z) \wedge \text{parent}(z, x))$$

Fact/Query: $\exists \vec{x} (p(\vec{t}_1) \wedge \dots \wedge p(\vec{t}_n)) = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$

Example:

$$F = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

Derivation: $F_0, (\rho_0, h_0, F_1), \dots, (\rho_n, h_n, F_{n+1})$

Example: Let $h(x) = \text{Joe}$.

$$1 \quad F_0 = \{\text{male}(\text{Joe}), \text{human}(\text{Joe})\}$$

$$2 \quad F_1 = \{\text{male}(\text{Joe}), \text{human}(\text{Joe}), \text{female}(z), \text{parent}(z, \text{Joe})\}$$

$$\mathcal{D} = F_0, (\rho, h, F_1).$$

A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules.
Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability

A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




fes

A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes

Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




A Jungle of Decidable Classes


Query Entailment: Let F be a fact and \mathcal{R} be a set of existential rules. Given query $Q = \{p(\vec{t}_1), \dots, p(\vec{t}_n)\}$, does $(F, \mathcal{R}) \models Q$?

Issue: Undecidable!

Solution: Restrict $\mathcal{R} \rightsquigarrow$ Decidability




Goal of This Project



Goals:

- ▶ Focus: BTS, Derivation Graphs, and Greediness
- ▶ Relationships b/w decidable classes
- ▶ Generalize existing notions


Goal of This Project



Goals:

- ▶ Focus: BTS, Derivation Graphs, and Greediness
- ▶ Relationships b/w decidable classes
- ▶ Generalize existing notions


Goal of This Project



Goals:

- ▶ Focus: BTS, Derivation Graphs, and Greediness
- ▶ Relationships b/w decidable classes
- ▶ Generalize existing notions

Goal of This Project



Goals:

- ▶ Focus: BTS, Derivation Graphs, and Greediness
- ▶ Relationships b/w decidable classes
- ▶ **Generalize existing notions**

- 1 Introduction
- 2 **BTS Notions**
- 3 Derivation Graphs
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion

Tree Decomposition

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- For each i , $X_i \subseteq \text{term}(F)$

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and


- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and

- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Tree Decomposition

Tree Decomposition: $T(F) = (V, E)$ with $V = \{X_1, \dots, X_n\}$ and


- ▶ For each i , $X_i \subseteq \text{term}(F)$
- ▶ $\bigcup_i X_i = \text{term}(F)$
- ▶ If $p(\vec{t}) \in F$, then $\exists X_i$ s.t. $\vec{t} \in X_i$
- ▶ $T(F) \upharpoonright t$ is connected

Example: Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$



Treewidth

Example (Cont.): Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Width: Largest node minus 1.

Treewidth: Smallest width among all tree decompositions.

Treewidth

Example (Cont.): Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Width: Largest node minus 1.

Treewidth: Smallest width among all tree decompositions.

Treewidth

Example (Cont.): Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Width: Largest node minus 1.

Treewidth: Smallest width among all tree decompositions.

Treewidth

Example (Cont.): Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$




Width: Largest node minus 1.

Treewidth: Smallest width among all tree decompositions.

Treewidth

Example (Cont.): Let $F = \{p(x, a), q(x, a, b), r(y, a, b)\}$



Width: Largest node minus 1.

Treewidth: Smallest width among all tree decompositions.

The Importance of Treewidth

Theorem (Courcelle, 1990)

The first-order theory of a class of first-order structures with bounded treewidth is decidable.

Bounded Treewidth Sets: A rule set \mathcal{R} is **bts** iff for any fact F , (F, \mathcal{R}) has a universal model of bounded treewidth.

Universal Model: A model for a FO theory that has a homomorphism into every other model.

Theorem (Baget et al. 2011)

*If \mathcal{R} is **bts**, then query entailment is decidable.*

Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.

Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.

Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.

Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.


Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Alternative Formulation of BTS: **adbts**

All-Derivation Bounded Treewidth Sets: \mathcal{R} is **adbts** iff for any F , there exists an $n \in \mathbb{N}$, if F' is derivable from F , then $tw(F') \leq n$.

Example: $F = \{p(a, b)\}$, $\mathcal{R} = \{p(x, y) \rightarrow p(y, z)\}$, $n = 1$.



Finding: **adbts** is a *proper* subset of **bts**.

Theorem

adbts \subsetneq **bts**

- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs**
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion

Derivation Graphs

Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Derivation Graphs

Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and

$\mathcal{R} = \{$


- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$

Derivation Graphs

Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and



$\mathcal{R} = \{$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$

Derivation Graphs


Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and

$\mathcal{R} = \{$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$



Derivation Graphs


Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and

$\mathcal{R} = \{$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$



Derivation Graphs


Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and

$\mathcal{R} = \{$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$



Derivation Graphs


Derivation Graph (Baget et al. 2011): A DAG that keeps track of how facts are derived.

Example: $F = \{p(x, y)\}$ and

$\mathcal{R} = \{$


- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z).$

$\}$




Derivation Graphs and Tree Decompositions

Idea (Baget et al. 2011): For a rule set \mathcal{R} , if derivation graph \rightsquigarrow tree for any fact, then \mathcal{R} is BTS.




Reduction Operations

Term Removal: If a term labels two converging arcs, it may be removed from one.




Reduction Operations

Arc Removal: If an arc is labelled with \emptyset , then it may be deleted.




Reduction Operations

Arc Contraction: If a node X sees a node Y and $terms(Y) \subseteq terms(X)$, then they can be merged.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.



Reduction Operations

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.



Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 *A reduced derivation graph is a tree decomposition.*
- 2 *Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.*

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 *A reduced derivation graph is a tree decomposition.*
- 2 *Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.*

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 *A reduced derivation graph is a tree decomposition.*
- 2 *Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.*

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 *A reduced derivation graph is a tree decomposition.*
- 2 *Width is bounded by $\max\{ |\text{terms}(F)|, |\text{terms}(\text{head}(R))|_{R \in \mathcal{R}} \}$.*

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 A reduced derivation graph is a tree decomposition.
- 2 Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 A reduced derivation graph is a tree decomposition.
- 2 Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Properties of Reduced Derivation Graphs


Theorem (Baget et al. 2011)

- 1 A reduced derivation graph is a tree decomposition.
- 2 Width is bounded by $\max\{ |terms(F)| , |terms(head(R))|_{R \in \mathcal{R}} \}$.

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

- 1 $p(x, y) \rightarrow q(y, z),$
- 2 $p(x, y) \wedge q(y, z) \rightarrow p(y, y),$
- 3 $p(y, y) \rightarrow q(y, y),$
- 4 $q(x, x) \wedge p(y, z) \rightarrow r(x, y, z)$

$$\}$$


Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

$$\blacktriangleright \text{adgs} \subset \text{wadgs} \subseteq \text{adbts} \subset \text{bts}$$

Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

$$\blacktriangleright \text{adgs} \subset \text{wadgs} \subseteq \text{adbts} \subset \text{bts}$$

Proof: Let \mathcal{R} be **wadgs** and F be a fact.

Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

$$\blacktriangleright \text{adgs} \subset \text{wadgs} \subseteq \text{adbts} \subset \text{bts}$$

Proof: Let \mathcal{R} be **wadgs** and F be a fact.

- 1 Assume F' is \mathcal{R} -derivable from F with \mathcal{D}

Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

$$\blacktriangleright \text{adgs} \subset \text{wadgs} \subseteq \text{adbts} \subset \text{bts}$$

Proof: Let \mathcal{R} be **wadgs** and F be a fact.

- 1 Assume F' is \mathcal{R} -derivable from F with \mathcal{D}
- 2 $DG(\mathcal{D})$ reducible to tree decomposition $T(F')$

Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

► **adgs** \subset **wadgs** \subseteq **adbts** \subset **bts**

Proof: Let \mathcal{R} be **wadgs** and F be a fact.

- 1 Assume F' is \mathcal{R} -derivable from F with \mathcal{D}
- 2 $DG(\mathcal{D})$ reducible to tree decomposition $T(F')$
- 3 $tw(F') \leq w(T(F')) \leq N = \max\{|terms(F)|, |terms(head(R))|_{R \in \mathcal{R}}\}$

Derivation Graph Sets

Acyclic Derivation Graph Set (adgs): A rule set \mathcal{R} is **adgs** iff for any F and any derivable F' , the derivation graph of every derivation of F' from F is reducible to a tree.

Weakly Acyclic Derivation Graph Set (wadgs): A rule set \mathcal{R} is **wadgs** iff for any F and any derivable F' , the derivation graph of some derivation of F' from F is reducible to a tree.

Theorem

- ▶ **adgs** \subset **wadgs** \subseteq **adbts** \subset **bts**
- ▶ *If \mathcal{R} is **adgs** or **wadgs**, then query entailment is decidable.*

Proof: Let \mathcal{R} be **wadgs** and F be a fact.

- 1 Assume F' is \mathcal{R} -derivable from F with \mathcal{D}
- 2 $DG(\mathcal{D})$ reducible to tree decomposition $T(F')$
- 3 $tw(F') \leq w(T(F')) \leq N = \max\{|terms(F)|, |terms(head(R))|_{R \in \mathcal{R}}\}$

- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs
- 4 Greedy Derivations**
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) Frontier of any rule application is (1) from the KB, or (2) from a single previous rule application.

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) **Frontier of any rule application** is (1) from the KB, or (2) from a single previous rule application.

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) Frontier of any rule application is (1) from the KB, or (2) from a single previous rule application.

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) Frontier of any rule application is (1) from the KB, or (2) from a single previous rule application.

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) Frontier of any rule application is (1) from the KB, or (2) from a single previous rule application.

Example:

$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x, y) \rightarrow q(y, z),$$

$$\mathbf{2} \quad p(x, y) \wedge q(y, z) \rightarrow q(y, x),$$

$$\mathbf{3} \quad q(y, x) \wedge q(y, z) \rightarrow r(x, y).$$

$$\}$$

Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) Frontier of any rule application is (1) from the KB, or (2) from a single previous rule application.

Example:


$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

$$1 \quad p(x, y) \rightarrow q(y, z),$$

$$2 \quad p(x, y) \wedge q(y, z) \rightarrow q(y, x),$$

$$3 \quad q(y, x) \wedge q(y, z) \rightarrow r(x, y).$$

$$\}$$


Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) **Frontier** of any rule application is (1) from the KB, or (2) from a single previous rule application.

Example:


$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

$$1 \quad p(x, y) \rightarrow q(y, z),$$

$$2 \quad p(x, y) \wedge q(y, z) \rightarrow q(y, x),$$

$$3 \quad q(y, x) \wedge q(y, z) \rightarrow r(x, y).$$

$$\}$$


Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) **Frontier** of any rule application is (1) from the KB, or (2) from a single previous rule application.

Example:


$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

$$1 \quad p(x, y) \rightarrow q(y, z),$$

$$2 \quad p(\textcolor{red}{x}, \textcolor{red}{y}) \wedge q(\textcolor{red}{y}, z) \rightarrow q(\textcolor{red}{y}, \textcolor{red}{x}),$$

$$3 \quad q(y, x) \wedge q(y, z) \rightarrow r(x, y).$$

$$\}$$


Greedy Derivations

Greedy Derivation (Thomazo et al. 2012) **Frontier** of any rule application is (1) from the KB, or (2) from a single previous rule application.

Example:


$$F = \{p(x, y)\}$$

$$\mathcal{R} = \{$$

$$1 \quad p(x, y) \rightarrow q(y, z),$$

$$2 \quad p(x, y) \wedge q(y, z) \rightarrow q(y, x),$$

$$3 \quad q(y, x) \wedge q(y, z) \rightarrow r(x, y).$$

$$\}$$


Greedy Derivations and Decidability

Greedy Bounded Treewidth Set (Thomazo et al. 2012): \mathcal{R} is **gbts** iff for all facts F and F' , if F' is derivable from F , then every derivation of F' from F is greedy.

Weakly Greedy Bounded Treewidth Set: \mathcal{R} is **wgbts** iff for all facts F and F' , if F' is derivable from F , then some derivation of F' from F is greedy.

Theorem

- ▶ **gbts** = **adgs**
- ▶ **gbts** \subset **wgbts** = **wadgs** \subseteq **bts**.
- ▶ If \mathcal{R} is **gbts** or **wgbts**, then query entailment is decidable.

Greedy Derivations and Decidability

Greedy Bounded Treewidth Set (Thomazo et al. 2012): \mathcal{R} is **gbts** iff for all facts F and F' , if F' is derivable from F , then every derivation of F' from F is greedy.


Weakly Greedy Bounded Treewidth Set: \mathcal{R} is **wgbts** iff for all facts F and F' , if F' is derivable from F , then some derivation of F' from F is greedy.

Theorem

- ▶ **gbts = adgs**
- ▶ **gbts** \subset **wgbts** = **wadgs** \subseteq **bts**.
- ▶ *If \mathcal{R} is **gbts** or **wgbts**, then query entailment is decidable.*


Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.




Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.



Greedy Derivations and Derivation Graphs

Cycle Removal: If there are arcs (X, Z) and (Y, Z) with an “ancestor” W such that $L(X, Z) \cup L(Y, Z) \subseteq \text{terms}(W)$, then (1) delete (X, Z) , (Y, Z) , (2) add (W, Z) , and (3) let $L(W, Z) = \text{Lab}(X, Z) \cup L(Y, Z)$.



Greedy Derivations and Decidability

Greedy Bounded Treewidth Set (Thomazo et al. 2012): \mathcal{R} is **gbts** iff for all facts F and F' , if F' is derivable from F , then every derivation of F' from F is greedy.

Weakly Greedy Bounded Treewidth Set: \mathcal{R} is **wgbts** iff for all facts F and F' , if F' is derivable from F , then some derivation of F' from F is greedy.

Theorem

- ▶ **gbts = adgs**
- ▶ **gbts** \subset **wgbts** = **wadgs** \subseteq **bts**.
- ▶ If \mathcal{R} is **gbts** or **wgbts**, then query entailment is decidable.

Greedy Derivations and Decidability

Greedy Bounded Treewidth Set (Thomazo et al. 2012): \mathcal{R} is **gbts** iff for all facts F and F' , if F' is derivable from F , then every derivation of F' from F is greedy.

Weakly Greedy Bounded Treewidth Set: \mathcal{R} is **wgbts** iff for all facts F and F' , if F' is derivable from F , then some derivation of F' from F is greedy.

Theorem

- ▶ **gbts** = **adgs**
- ▶ **gbts** \subset **wgbts** = **wadgs** \subseteq **bts**.
- ▶ If \mathcal{R} is **gbts** or **wgbts**, then query entailment is decidable.

- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts****
- 6 Conclusion

Rule Dependencies

Dependency (Baget 2004): A rule ρ' depends on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$\mathcal{R} = \{$

1 $p(x, y) \rightarrow q(x, y)$

2 $q(x, y) \rightarrow r(x, y)$

$\}$

Rule Dependencies

Dependency (Baget 2004): A rule ρ' **depends** on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$\mathcal{R} = \{$

- 1 $p(x, y) \rightarrow q(x, y)$

- 2 $q(x, y) \rightarrow r(x, y)$

$\}$

Rule Dependencies

Dependency (Baget 2004): A rule ρ' depends on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$$\mathcal{R} = \left\{ \begin{array}{ll} \text{1} & p(x, y) \rightarrow q(x, y) \\ \text{2} & q(x, y) \rightarrow r(x, y) \end{array} \right. \quad \text{1} \quad \{p(a, b)\}$$

Rule Dependencies

Dependency (Baget 2004): A rule ρ' depends on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$\mathcal{R} = \{$

1 $p(x, y) \rightarrow q(x, y)$

2 $q(x, y) \rightarrow r(x, y)$

$\}$

1 $\{p(a, b)\}$

2 $\{p(a, b), q(a, b)\}$

Rule Dependencies

Dependency (Baget 2004): A rule ρ' depends on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$\mathcal{R} = \{$

1 $p(x, y) \rightarrow q(x, y)$

2 $q(x, y) \rightarrow r(x, y)$

$\}$

1 $\{p(a, b)\}$

2 $\{p(a, b), q(a, b)\}$

3 $\{p(a, b), q(a, b), r(a, b)\}$

Rule Dependencies

Dependency (Baget 2004): A rule ρ' depends on a rule ρ , if there exists a fact F such that applying ρ to F gives a fact F' that ρ' can be applied to (and was not previously applicable to).

Example: Let $F = \{p(a, b)\}$

$$\mathcal{R} = \left\{ \begin{array}{ll} \text{1} & p(x, y) \rightarrow q(x, y) \\ \text{2} & q(x, y) \rightarrow r(x, y) \end{array} \right. \quad \begin{array}{l} \text{1} \quad \{p(a, b)\} \\ \text{2} \quad \{p(a, b), q(a, b)\} \\ \text{3} \quad \{p(a, b), q(a, b), r(a, b)\} \end{array}$$

$q(x, y) \rightarrow r(x, y)$ depends on $p(x, y) \rightarrow q(x, y)$

Graph of Rule Dependencies

Graph of Rule Dependencies (Baget 2004): A directed graph showing all dependencies between rules.

$\mathcal{R} = \{$

1 $\rho_1 = p(x) \rightarrow q(x, y)$

2 $\rho_2 = r(x) \rightarrow s(x, y, z)$

3 $\rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $\rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$

Graph of Rule Dependencies

Graph of Rule Dependencies (Baget 2004): A directed graph showing all dependencies between rules.

$\mathcal{R} = \{$

1 $\rho_1 = p(x) \rightarrow q(x, y)$

2 $\rho_2 = r(x) \rightarrow s(x, y, z)$

3 $\rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $\rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Graph of Rule Dependencies

Graph of Rule Dependencies (Baget 2004): A directed graph showing all dependencies between rules.

$\mathcal{R} = \{$


1 $\rho_1 = p(x) \rightarrow q(x, y)$

2 $\rho_2 = r(x) \rightarrow s(x, y, z)$

3 $\rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $\rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Graph of Rule Dependencies

Graph of Rule Dependencies (Baget 2004): A directed graph showing all dependencies between rules.

$\mathcal{R} = \{$


1 $\rho_1 = p(x) \rightarrow q(x, y)$

2 $\rho_2 = r(x) \rightarrow s(x, y, z)$

3 $\rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $\rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Graph of Rule Dependencies

Graph of Rule Dependencies (Baget 2004): A directed graph showing all dependencies between rules.

$\mathcal{R} = \{$


1 $\rho_1 = p(x) \rightarrow q(x, y)$

2 $\rho_2 = r(x) \rightarrow s(x, y, z)$

3 $\rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $\rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



An Observation


 $\mathcal{R} = \{$

$$1 \quad \rho_1 = p(x) \rightarrow q(x, y)$$

$$2 \quad \rho_2 = r(x) \rightarrow s(x, y, z)$$

$$3 \quad \rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$4 \quad \rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

 $\}$


A Derivation: Let $\rho \in \{\rho_1, \rho_2, \rho_3\}$.

$$F_0, \dots, (\rho_i, h_i, F_i), (\rho, h_{i+1}, F_{i+1}), \dots, (\rho_n, h_n, F_n)$$

An Observation


 $\mathcal{R} = \{$

$$1 \quad \rho_1 = p(x) \rightarrow q(x, y)$$

$$2 \quad \rho_2 = r(x) \rightarrow s(x, y, z)$$

$$3 \quad \rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$4 \quad \rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

 $\}$


A Derivation: Let $\rho \in \{\rho_1, \rho_2, \rho_3\}$.

$$F_0, \dots, (\rho_i, h_i, F_i), (\rho, h_{i+1}, F_{i+1}), \dots, (\rho_n, h_n, F_n)$$

An Observation


 $\mathcal{R} = \{$

$$\mathbf{1} \quad \rho_1 = p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad \rho_2 = r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad \rho_3 = p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad \rho_4 = q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

 $\}$


A Derivation: Let $\rho \in \{\rho_1, \rho_2, \rho_3\}$.

$$F_0, \dots, (\rho_i, h_i, F_i), (\rho, h_{i+1}, F_{i+1}), \dots, (\rho_n, h_n, F_n)$$

The Observation: We can permute rule applications!

$$F_0, \dots, (\rho, h_{i+1}, F_{i-1} \cup (F_{i+1} \setminus F_i)), (\rho_i, h_i, F_{i+1}), \dots, (\rho_n, h_n, F_n)$$

Re-writing Derivations via Permutations

Lemma (Permutation Lemma)

Suppose we have a derivation of the following form:

$$\mathcal{D} := F_0, \dots, (\rho_i, h_i, F_{i+1}), (\rho_{i+1}, h_{i+1}, F_{i+2}), \dots, (\rho_{n-1}, h_{n-1}, F_n)$$

If ρ_{i+1} does not depend on ρ_i , then the following is a derivation as well:

$$\mathcal{D}' := F_0, \dots, (\rho_{i+1}, h_{i+1}, F'_i), (\rho_i, h_i, F_{i+2}), \dots, (\rho_{n-1}, h_{n-1}, F_n)$$

where $F'_i = F_i \cup (F_{i+2} - F_{i+1})$.

A Useful Rule Set for Non-Greediness

$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$


$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$

A Useful Rule Set for Non-Greediness

 $\mathcal{R} = \{$

1 $p(x) \rightarrow q(x, y)$

2 $r(x) \rightarrow s(x, y, z)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

 $\}$

A Useful Rule Set for Non-Greediness


 $\mathcal{R} = \{$

1 $p(x) \rightarrow q(x, y)$

2 $r(x) \rightarrow s(x, y, z)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

 $\}$ 

A Useful Rule Set for Non-Greediness


 $\mathcal{R} = \{$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

 $\}$


A Useful Rule Set for Non-Greediness


 $\mathcal{R} = \{$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

 $\}$


A Useful Rule Set for Non-Greediness


$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$


\mathcal{R} is not greedy!

But is \mathcal{R} Weakly Greedy?!

$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$


$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$

But is \mathcal{R} Weakly Greedy?!


$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$

But is \mathcal{R} Weakly Greedy?!


$$\mathcal{R} = \{$$

$$1 \quad p(x) \rightarrow q(x, y)$$

$$2 \quad r(x) \rightarrow s(x, y, z)$$

$$3 \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$4 \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$


But is \mathcal{R} Weakly Greedy?!


$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$


But is \mathcal{R} Weakly Greedy?!


$$\mathcal{R} = \{$$

$$\mathbf{1} \quad p(x) \rightarrow q(x, y)$$

$$\mathbf{2} \quad r(x) \rightarrow s(x, y, z)$$

$$\mathbf{3} \quad p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$$

$$\mathbf{4} \quad q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$$

$$\}$$


Perhaps \mathcal{R} is weakly greedy!

Showing \mathcal{R} is weakly greedy

Idea 1: Use 3 to simulate 1 and 2.

$\mathcal{R} = \{$


1 $p(x) \rightarrow q(x, y)$

2 $r(y) \rightarrow s(y, u, v)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Showing \mathcal{R} is weakly greedy

Idea 1: Use 3 to simulate 1 and 2.

$\mathcal{R} = \{$


1 $p(x) \rightarrow q(x, y)$

2 $r(y) \rightarrow s(y, u, v)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Showing \mathcal{R} is weakly greedy

Idea 1: Use 3 to simulate 1 and 2.

$\mathcal{R} = \{$


1 $p(x) \rightarrow q(x, y)$

2 $r(y) \rightarrow s(y, u, v)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Showing \mathcal{R} is weakly greedy

Idea 1: Use 3 to simulate 1 and 2.

$\mathcal{R} = \{$


1 $p(x) \rightarrow q(x, y)$

2 $r(y) \rightarrow s(y, u, v)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$



Showing \mathcal{R} is weakly greedy

Idea 1: Use 3 to simulate 1 and 2.

$\mathcal{R} = \{$


1 $p(x) \rightarrow q(x, y)$

2 $r(y) \rightarrow s(y, u, v)$

3 $p(x) \wedge r(y) \rightarrow q(x, z) \wedge s(y, u, v)$

4 $q(x, y) \wedge s(z, u, v) \rightarrow t(x, z, u, w)$

$\}$




Idea 2: Permutation Lemma \rightsquigarrow

- (i) Permute instances of 1 or 2 backward to instances of 2 or 1
- (ii) Replace 1,2 or 2,1 instances with 3.

- 1 Introduction
- 2 BTS Notions
- 3 Derivation Graphs
- 4 Greedy Derivations
- 5 Showing **gbts** \subset **wgbts**
- 6 Conclusion**

Conclusion



*Note: Proof must be checked in detail.