

Decidability of Quasi-Dense Modal Logics

Tim Lyon¹ and Piotr Ostropolski-Nalewaja^{1,2}

¹TU Dresden ²University of Wrocław



Table of contents

1. Introduction
2. Our paper
3. A glimpse into the proof

Introduction

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

$$\text{Expression} \quad k \rightarrow k'$$

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{k \rightarrow k'\}$ enforces that in models of φ every pair of nodes u, v satisfy:

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{ k \rightarrow k' \}$ enforces that in models of φ every pair of nodes u, v satisfy:

If u and v are connected with a path of length k ...

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{ k \rightarrow k' \}$ enforces that in models of φ every pair of nodes u, v satisfy:

*If u and v are connected with a path of length k ...
...then $they$ are connected with a path of length k' .*

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{k \rightarrow k'\}$ enforces that in models of φ every pair of nodes u, v satisfy:

*If u and v are connected with a path of length k ...
...then $they$ are connected with a path of length k' .*

$2 \rightarrow 1$ is *transitivity*

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{k \rightarrow k'\}$ enforces that in models of φ every pair of nodes u, v satisfy:

*If u and v are connected with a path of length k ...
...then $they$ are connected with a path of length k' .*

$2 \rightarrow 1$ is *transitivity* $0 \rightarrow 1$ is *reflexivity*

Setting up the stage

Considered logic

$$\varphi ::= p \mid \neg p \mid \varphi \vee \varphi \mid \varphi \wedge \varphi \mid \Diamond \varphi \mid \Box \varphi$$

Path axioms / modal reduction principles

Expression $k \rightarrow k'$ where $k, k' \in \mathbb{N}$

Also denoted as $\Diamond^k p \rightarrow \Diamond^{k'} p$

$\{ k \rightarrow k' \}$ enforces that in models of φ every pair of nodes u, v satisfy:

*If u and v are connected with a path of length k ...
...then $they$ are connected with a path of length k' .*

$2 \rightarrow 1$ is *transitivity* $0 \rightarrow 1$ is *reflexivity* $1 \rightarrow 2$ is *density*

We are interested in deciding satisfiability of extensions of K with MRPs:

Open and closed cases

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

Open and closed cases

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

Open and closed cases

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

transitivity ($2 \rightarrow 1$)

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

transitivity ($2 \rightarrow 1$)

reflexivity + transitivity

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

transitivity ($2 \rightarrow 1$)

reflexivity + transitivity

density ($1 \rightarrow 2$)

See e.g. P. Blackburn, M. de Rijke, Y. Venema. *Modal Logic* (2001).

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

transitivity ($2 \rightarrow 1$)

reflexivity + transitivity

density ($1 \rightarrow 2$)

See e.g. P. Blackburn, M. de Rijke, Y. Venema. *Modal Logic* (2001).

General transitivity ($k \rightarrow 1$)

via a standard filtration argument.

Open and closed cases

We are interested in deciding satisfiability of extensions of K with MRPs:

General case ($k \rightarrow k'$) is open

J. van Benthem. *Modal reduction principles* (1976).

reflexivity ($0 \rightarrow 1$)

transitivity ($2 \rightarrow 1$)

reflexivity + transitivity

density ($1 \rightarrow 2$)

See e.g. P. Blackburn, M. de Rijke, Y. Venema. *Modal Logic* (2001).

General transitivity ($k \rightarrow 1$)

via a standard filtration argument.

K4 with any MRPs

M. Zakharyashev. Canonical formulas for K4. Part I: Basic results (1992).

Our paper

Quasi-Dense Modal Reduction Principle

is any MRP of the form $k \rightarrow k_+$ where $k < k_+$.

Quasi-Dense Modal Reduction Principle

is any MRP of the form $k \rightarrow k_+$ where $k < k_+$.

Quasi-Dense Modal Logic

is any extension of K with a finite number of Quasi-Dense MRPs.

In this paper

In this paper

**Quasi-Dense Modal Logics can be decided in
EXPSPACE.**

A glimpse into the proof

Step I: RE-Model Building Procedure

Step I: RE-Model Building Procedure

Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with $\bullet P_\phi$

Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ

For $\Diamond\psi \subseteq \varphi$



Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ

For $\Diamond\psi \subseteq \varphi$



For $\Box\psi \subseteq \varphi$



Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ

For $\Diamond\psi \subseteq \varphi$



For $\Box\psi \subseteq \varphi$



For $\psi \vee \psi' \subseteq \varphi$



Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ

For $\Diamond\psi \subseteq \varphi$



For $\Box\psi \subseteq \varphi$



For $\psi \vee \psi' \subseteq \varphi$



For $\psi \wedge \psi' \subseteq \varphi$



Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ


For $\Diamond\psi \subseteq \varphi$



For $\Box\psi \subseteq \varphi$



For $\psi \vee \psi' \subseteq \varphi$



For $\psi \wedge \psi' \subseteq \varphi$




For $k \rightarrow k_+ \in P$



Step I: RE-Model Building Procedure

For $\varphi \in K$ and set P of QDMRPs.

Start with P_ϕ



For $\Diamond\psi \subseteq \varphi$



For $\Box\psi \subseteq \varphi$



For $\psi \vee \psi' \subseteq \varphi$



For $\psi \wedge \psi' \subseteq \varphi$



For $k \rightarrow k_+ \in P$



If P_p and $P_{\neg p}$ appear over a node

derive nullary predicate contr.

We have that:

$\langle \varphi, P \rangle$ is satisfiable *iff* The procedure non-deterministically derives a structure that does not contain contr.

A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$

$$\begin{array}{c} \bullet \\ \Box\Box p \end{array}$$

A quick example

$$\varphi = \Box\Box p$$

$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$


$$P = \{2 \rightarrow 3\}$$



A quick example

$$\varphi = \Box\Box p$$

$$P = \{2 \rightarrow 3\}$$



A more complex example

$$\varphi = \Box\Box p \wedge \Diamond\Diamond(p \wedge \Diamond\neg p \quad \vee \quad \neg p \wedge \Diamond p)$$

$$P = \{2 \rightarrow 3\}$$

$$\begin{array}{c} \bullet \\ \Box\Box p \end{array}$$

A more complex example

$$\varphi = \Box\Box p \wedge \Diamond\Diamond(p \wedge \Diamond\neg p \vee \neg p \wedge \Diamond p)$$


$$P = \{2 \rightarrow 3\}$$



A more complex example

$$\varphi = \Box\Box p \wedge \Diamond\Diamond(p \wedge \Diamond\neg p \vee \neg p \wedge \Diamond p)$$


$$P = \{2 \rightarrow 3\}$$



A more complex example

$$\varphi = \Box\Box p \wedge \Diamond\Diamond(p \wedge \Diamond\neg p \quad \vee \quad \neg p \wedge \Diamond p)$$

$$P = \{2 \rightarrow 3\}$$



We start with the set S of all structures that can be derived by the non-deterministic model-building procedure.

We start with the set \mathcal{S} of all structures that can be derived by the non-deterministic model-building procedure.

We find a way to represent set \mathcal{S} via a finite set of bounded-size structures called *templates*.

We start with the set \mathcal{S} of all structures that can be derived by the non-deterministic model-building procedure.

We find a way to represent set \mathcal{S} via a finite set of bounded-size structures called *templates*.

We enumerate the templates, and since their size is at most exponential in $\langle \varphi, P \rangle$, we get an EXPSPACE algorithm for checking satisfiability.

We start with the set \mathcal{S} of all structures that can be derived by the non-deterministic model-building procedure.

We find a way to represent set \mathcal{S} via a finite set of bounded-size structures called *templates*.

We enumerate the templates, and since their size is at most exponential in $\langle \varphi, P \rangle$, we get an EXPSPACE algorithm for checking satisfiability.

Thank you!

Questions?